

# Restoring anterior aesthetics with two-piece zirconia implants

Dr Saurabh Gupta, India

Zirconia implants are one of the newest and most exciting developments in dental implantology. Multiple studies have proved that zirconia implants induce little to no peri-implant tissue inflammation and allow for high levels of epithelial attachment. Additionally, these implants look more natural; hence, they provide improved aesthetics. Furthermore, they do not have metal components, which makes them ideal for people with metal sensitivities and patients who would prefer their implants to be metal-free.<sup>1-3</sup>

Aesthetics around natural teeth can be challenging under normal circumstances. When teeth are to be replaced with implants, especially in the aesthetic zone, gingival tissue can complicate the desired results. In a patient with a thin gingival biotype, the grey of a titanium implant will show through, leading to a darker gingiva overlying that area and decreasing the aesthetics of the patient's smile. A patient who has had a missing anterior tooth for a period, resulting in resorption of the facial plate even with a thicker gingival biotype, will have less bone over the implant on the facial aspect of the ridge. The result, like with a thin gingival biotype, will be a shadow over the underlying implant that hampers the aesthetic result and does not blend with the adjacent tissue around the natural teeth.<sup>4-7</sup> In the following, a clinical case is described to demonstrate the use of two-piece ceramic implants in the anterior aesthetic zone to avoid this difficulty.

## Case presentation

A 44-year-old male patient presented to our office to learn about options for replacement of his failing maxillary central incisors after undergoing partial root canal therapy. He also complained of greyish gingiva around the endodontically treated teeth and desired a metal-free solution (Figs. 1 & 2). Photographs of his teeth when smiling were taken to assess the overall aesthetic risk of the case. Treatment options were then discussed with the patient. After reviewing the options, the patient chose to have the endodontically treated teeth extracted and replaced with two-piece zirconia implants and metal-free crowns.

## Surgical procedure

The guidelines for zirconia implant placement in the anterior zone and the drilling protocol specified by the manufacturer (Zeramex XT, Dentalpoint) were followed. It is important to note that implant sites must be prepared adequately to prevent excessive implant insertion torque and that the use of a bone tap is necessary. Both the



Fig. 1: Initial situation. Fig. 2: Initial radiograph.

# ISMI

# INT. SOCIETY OF METAL FREE IMPLANTOLOGY



The innovative specialist society for modern metal-free implantology

## Become a member now!

### Benefits



#### Effective public relations

Benefit from a strong community that elevates the marketing of your practice through effective public relations strategies.

#### Personal online member profile

ISMI provides a personal profile of all active members on their website—free of charge. In addition, the ISMI patient platform provides important information for patients and features a search tool with which patients can find their perfect dentist.



#### Discount on congress fees

Come and join us! Attend the Annual Meeting of ISMI in Hamburg on 3 and 4 May 2024 and benefit from first-rate continuing education. ISMI members receive a special discount on the participation fee.

#### Online archive for specialists

Get exclusive access to ISMI's extensive online archive. Discuss all relevant questions regarding metal-free implantology with experts and colleagues from around the world and enjoy free access to the online archive where you will find informative training videos and clinical case reports.



#### Newsletter

The ISMI newsletter keeps you up to date with the latest scientific trends, products, and events on a regular basis. It also features user reports as well as a wide range of information and tips on the subject of metal-free implantology.

#### Specialist magazine


As a member of ISMI, your membership fee includes a subscription of the independently published English language magazine *ceramic implants*—international magazine of ceramic implant technology. Published three times per year, the magazine offers specialist articles and event reports as well as industry- and science-related news from the international world of metal-free implantology. In addition, *ceramic implants* provides information about manufacturers and their latest products.



ISMI e.V. Office Leipzig  
Holbeinstraße 29 | 04229 Leipzig | Germany  
Phone: +49 800 4764-000 | Fax: +49 800 4764-100  
office@ismi.me | www.ismi.me

Become a  
member of ISMI now!





**Fig. 3:** After atraumatic extraction of teeth #11 and 21. **Fig. 4:** Immediate placement of Zeramex XT implants. **Fig. 5:** PMMA temporary crowns placed one week after surgery.

vertical and transverse insertion depth of the zirconia implant are important for prosthetic success. The implant can be placed between 1.6 and 0.6mm supra-crestally because the neck section (0.6mm) is smooth. The insertion depth is determined by the height of the gingiva and the existing bone around the adjacent teeth.

After atraumatic extractions and laser curettage, two-piece zirconia implants (4.2×12.0mm) were placed in sites #11 and 21 under local anaesthesia, cover screws were placed and the sites closed to allow for healing (Figs. 3 & 4). After 72 hours, the PMMA temporary crowns were inserted (Fig. 5).

After a four-month healing period, the second-stage surgery was performed with a 940nm diode laser, the cover screws were removed and healing abutments were placed for a period of two weeks. Reduced inflammation of the peri-implant soft tissue was noted, demonstrating excellent biocompatibility and host response (Fig. 6).

Angulated abutments (15°), also made of alumina-toughened zirconia like the implants, were placed on the implants with Zeramex XT VICARBO screws (Figs. 7–9). This screw, which is made of longitudinal carbon fibre strands and moulded slightly larger than the internal aspect of the implant, allows absorption of the forces of mastication.



**Fig. 8:** Angulated zirconia abutments in position. **Fig. 9:** Four-month post-op radiograph. **Fig. 10:** Zirconia crowns *in situ*, lateral view. **Fig. 11:** Zirconia crowns *in situ*, frontal view.


## Specialist in Integrative Oral Medicine

- Become a member of **ICOSIM e.V.** & get access to the **IOM-Curriculum**

- Access to over **6 learning blocks** with more than **40 learning videos**

- Free downloads with **exclusive bonus material**

- Participation in **FDOK-Weeks inkl. Live-OPs & face-to-face training**



**Fig. 6:** Healing after four months. **Fig. 7:** Example of a Zeramex XT implant, abutment and VICARBO screw (metal-free solution).


tion and provides a hermetically sealed connection. A digital impression was taken for the fabrication of the final crowns (Fig. 10). Zirconia crowns were cemented to the abutment heads with glass ionomer cement to provide natural aesthetics. Instructions were given for maintenance and periodic recall (Figs. 10–12).

### Discussion

Owing to rising complications observed in some clinical situations involving the use of titanium dental implants and the growing incidence of peri-implant mucositis and peri-implantitis affecting both the short- and long-term survival rates of titanium dental implants, the develop-

ment of alternative materials to address these has been pursued. Zirconia has been shown to have similar osseointegration success to titanium, offer a soft-tissue response that is superior to that of titanium and have less of an affinity for plaque collection compared with titanium surfaces.


Also, the peri-implant soft tissue around titanium and zirconia abutments has been shown to have colour differences compared with the soft tissue around natural teeth, and the peri-implant soft tissue around zirconia has been demonstrated to have a better colour match to the soft tissue than titanium. This can be extrapolated to the aesthetics of the colour of the implant itself. Zirconia implants can be used in aesthetic situations, the white shade of the implant eliminating any potential for darkening of the gingival tissue and providing a more natural final aesthetic result than is possible with titanium implants. Long-term studies are necessary to continue to evaluate the effectiveness and success rates of two-piece zirconia implants.



**Fig. 12:** Radiograph of implants and final crowns.

### contact

**Dr Saurabh Gupta**  
+91 9916 203455  
saurabh@iaoci.com



**Now**  
Become a  
member of  
**ICOSIM e.V.** &  
register for the  
**IOM-curriculum:**



**More information:**  
[www.icosim.de](http://www.icosim.de)