

Full arch reconstruction of the edentulous maxilla

From bone augmentation to final monolithic zirconia restoration

Prof. João Caramês, Portugal

Dental implant rehabilitation in young patients with severe alveolar bone loss poses a significant challenge, particularly when previous implant attempts have failed. This case report describes the treatment of a 20-year-old female patient who had previously undergone unsuccessful implant treatment and was reluctant to consider removable dentures as a permanent solution.



Figs. 1a–c: Missing maxillary teeth causing face flattening in a young patient.




Fig. 2: Radiograph assessment showed limited bone availability.

The treatment plan focused on restoring function and aesthetics through a combination of bone augmentation, sinus lift, and implant placement using Straumann® Bone Level Tapered (BLT) implants. The surgical workflow included the use of Straumann® Xenograft for ridge augmentation, followed by digital planning for guided implant surgery. A temporary removable denture was provided during the healing phase, after which the patient received a screw-retained monolithic zirconia prosthesis.

The patient-centered approach prioritised the young patient's well-being and limited bone anatomy. A two-stage strategy was chosen to optimise conditions for standard implants, ensuring stable, long-term outcomes. After reviewing treatment options and their pros and cons, the patient selected this approach.




Fig. 3: Based on SAC classification, the case was categorised as complex.





Fig. 4: Full-thickness flap with crestal incision for surgical site exposure. – **Fig. 5:** Bone augmentation was done using Straumann® Xenograft®. – **Fig. 6:** After one month, soft tissue heals with no complications.

Beyond the clinical success of the procedure, the impact on the patient's quality of life was evident. Prior to treatment, the patient reported considerable difficulties with masticatory function, poor nutrition, and a decline in self-esteem due to the limitations of removable prosthetics. Following the full-arch rehabilitation, the patient experienced marked improvements in oral function, aesthetics, and psychosocial well-being. This case highlights the critical role that implant-supported restorations play in not only addressing dental health issues but also enhancing overall patient satisfaction and quality of life.

Initial situation

A 20-year-old woman came to our clinic in good health, not taking any medication, and with no allergies. She told us that she had lost her teeth at a young age because of cavities. Although she had tried implants before, the treatment did not work out, and she could not imagine wearing removable dentures for the rest of her life. She wanted to find a fix solution using dental implants to restore her failing teeth.

The extra-oral examination showed a flattening of the lower third of the face, attributed to the lack of support from the missing maxillary teeth (Figs. 1a–c).

The intra-oral examination revealed an edentulous maxilla characterised by significant horizontal and vertical bone resorption. This considerable loss of bone structure presents substantial challenges for implant placement, as it may compromise the necessary stability and support for a successful prosthesis.

Additionally, the radiographic examination confirmed recent implant loss and demonstrated a critical deficiency of bone in both vertical and horizontal dimensions (Fig. 2). This imaging underscores the necessity for thorough evaluation and intervention to address the underlying bone deficiencies, highlighting the urgent need for a comprehensive treatment plan to effectively restore both function and aesthetics.

Based on the SAC classification, the patient's surgical and prosthodontic case was categorised as complex (Fig. 3).

Treatment planning

The treatment plan was tailored to meet the patient's functional and aesthetic needs while ensuring long-term stability. It followed a structured approach, beginning with sinus lift and bone augmentation and digital planning, continuing through implant placement, and ending with the delivery of a final

screw-retained prosthesis. Each step was carefully planned to achieve a successful outcome.

The treatment workflow included:

1. Maxilla reconstruction: Bilateral sinus lift, vertical and horizontal bone augmentation with Straumann® Xenograft.
2. Temporary removable denture during the six-month healing period.
3. Digital planning for guided implant surgery and immediate loading protocol.
4. Insertion of eight Straumann® Bone Level Tapered SLActive® implants.

5. Delivery of screw-retained immediate prosthesis.
6. Monitoring during osseointegration period.
7. Finalisation with a screw-retained zirconia monolithic prosthesis.

The decision was guided by the doctor's patient-centered approach, which emphasised the patient's overall well-being. Given the patient's young age and limited bone anatomy unsuitable for standard implants, the doctor adopted a thoughtful two-stage strategy. This approach, tailored to the patient's age, preferences, and healthy systemic condition, aims to enhance the anatomical foundation for standard implant placement, ensuring stable, predictable, long-term outcomes. Comprehensive information about various treatment options, including their benefits and drawbacks, was provided to the patient, who ultimately chose the current approach as her preferred treatment plan.

Surgical procedure

Local anaesthesia with lidocaine 2% with epinephrine 1:100,000 was administered. The surgical procedure began with a full-thickness flap created through a crestal incision to expose the surgical site (Fig. 4). Bone augmentation was performed using Straumann® Xenograft® to enhance the alveolar ridge and provide adequate support for future implants (Fig. 5).

One month after the procedure, the soft tissues exhibited good healing progress, showing a healthy appearance with no signs of complications (Fig. 6). This positive outcome reflects the effectiveness of the surgical technique and the patient's compliance to postoperative care instructions.

At the six-month follow-up, a CBCT scan revealed that the augmented bone had fully integrated, providing the necessary dimensions and stability for implant placement (Fig. 7). Using digital planning software, the 3D positioning of the implants was meticulously planned to ensure optimal placement (Fig. 8). This




Fig. 7: Six-month CBCT confirms successful bone integration for implant placement. – **Fig. 8:** Precise implant positioning with 3D digital planning.





Fig. 9: Surgical site preparation for implant insertion. – **Fig. 10:** Optimal implant placement with guided surgery.

meticulous planning ensures accurate angulation and depth of the implants, significantly improving both functional performance and aesthetic outcomes of the final prosthesis.

During the second surgery, the surgical guide was checked for proper fit. A full-thickness flap was again raised, the fixation pins were then drilled and inserted, and the implants were inserted using a fully guided technique to maximise accuracy (Fig. 9). Straumann® BLT implants were placed with precision aid of the handpiece in a clockwise direction with a speed of 15rpm and torqued at 35Ncm (Fig. 10).

Prosthetic procedure

A temporary screw-retained prosthesis was delivered immediately, following the immediate loading protocol, to maintain function and aesthetics during the healing phase (Fig. 11). Oral hygiene instructions were given, and occlusion checked.

At the suture removal appointment, healing was observed to be uneventful. The patient was closely monitored throughout the osseointegration period, and treatment was finalised with the placement of a screw-retained monolithic zirconia prosthesis

Treatment outcomes

The final outcome of the treatment emphasises the aesthetic excellence of the zirconia monolithic prosthesis (Fig. 12), which blends seamlessly with the natural dentition. Its functional durability ensures long-term stability, providing both visual appeal and enhanced performance. This result has greatly improved the patient's overall satisfaction and quality of life.

Additionally, lateral views taken at the end of the treatment offer a thorough perspective on the placement and fit of the prosthesis. These images confirm



Fig. 11: Immediate loading with temporary screw-retained prosthesis. – **Fig. 12:** Final restoration with zirconia monolithic prosthesis.

proper alignment with the surrounding tissues and occlusion, ensuring both functional performance and aesthetic harmony, further contributing to the overall success of the treatment (Figs. 13a–c).

A radiographic control performed one-year post-surgery confirms the successful integration and stability of the implants, providing further evidence of the long-term success of the procedure (Fig. 14). This imaging not only highlights the osseointegration achieved but also reassures the reliability of the implants in supporting the prosthesis, contributing to the patient's overall dental health.

The final outcome demonstrates the excellent health of both hard and soft tissues. The patient expressed great grati-

tude and satisfaction, saying she could now smile and be herself again, after what had once been a nightmare.

Author's testimonial

Critical risk factors should be carefully evaluated prior to deciding the loading protocol. Different levels of bone atrophy should require different rehabilitation approaches. The Carames Classification system for full arch implant rehabilitation can support you in the choice of the optimal treatment. Regenerative procedures provide an added value for the patient's present and future rehabilitation needs. Cross-arch stabilisation is a key factor in full-arch immediate loading.

About...

Prof. João Caramês pursued advanced expertise in dentistry at New York University, earning a postgraduate degree in oral rehabilitation and implantology and becoming the first non-American to receive its prestigious Alumni Award.

Back in Portugal, he founded the Instituto de Implantologia®, revolutionising dentistry with cutting-edge knowledge acquired abroad.

Currently, he is a guest professor and International director for NYU's continuing education department and leads the implantology research line at the Oral and Biomedical Sciences Research Unit (UICOB). With over 30 years of experience and more than 50,000 implants placed, he is a global authority in implantology and the creator of the widely adopted "Caramês Classification", a system tailored to individual patient needs.



Figs. 13a–c: One year after final restoration, the patient was highly satisfied with stable results.




Fig. 14: One-year radiograph confirms implant success and stability.

Author details



Contact address

Prof. João Caramês

Instituto de Implantologia

Lisbon, Portugal

md@institutodeimplantologia.pt

www.institutodeimplantologia.pt



ZWP ONLINE


www.zwp-online.info




© Kelly-Sikkema - unsplash.com

show some

leave a like.

 [facebook.com/
zwponline](https://facebook.com/zwponline)

 [instagram.com/
zwponline](https://instagram.com/zwponline)

 [x.com/
zwponline](https://x.com/zwponline)

