

20th European Consensus Conference (EuCC) updates its guideline

EuCC recommendations on complications in implant treatment

The aim of the 2025 guideline of the 20th European Consensus Conference (EuCC) under the auspices of BDIZ EDI is to prevent complications and, if they do occur, to treat them correctly in order to improve the patient outcome. The new eight-page paper is intended to serve as a recommendation for implantologists on how to deal with complications.



The international panel of 16 experts at the European Consensus Conference discussed the various treatment concepts in implant therapy, including both surgical and prosthetic approaches.

They concluded: "Dental implants are reliable treatment options for restoring patient function and aesthetics. Careful

case selection is necessary by considering not only the oral findings alone. Due to the great variation of implant designs and surgical and prosthetic procedures proposed, the individual suggested parameter should be followed to avoid

complications. All procedures should be performed by treatment providers with the requisite up-to-date expertise and training.

The working paper was prepared by Prof. Jörg Neugebauer of the Department of Oral, Maxillofacial and Plastic Facial Surgery, Polyclinic for Oral Surgery and Implantology, Plastic, Reconstructive and Aesthetic Surgery, University of Cologne, Germany. The members of the EuCC reviewed and discussed the first draft according to the following timetable: review of the first draft, registration of alternative proposals, voting on recommendations and levels of recommendation, discussion of points of disagreement, and final voting.

Changes from the 2019 version

As the host of the European Consensus Conference, Neugebauer pointed out the changes to the previous 2019 version: "It was important for this year's EuCC to highlight the need to develop specific, patient-centred treatment strategies. The restrictive approach to osteoporosis is also outdated. A 2023 meta-analysis by Lemos CAA et al. has prompted the EuCC to issue the cautiously optimistic statement that there is 'no increased general risk of osteoporosis'."

Ordering the guideline

The 8-page guideline 2025 brochure with its comprehensive bibliography in German or English can be ordered from the BDIZ EDI online shop at a price of €4.50 per copy (including VAT, plus shipping).

A sample is also available in the online shop.

Members will receive the guideline free of charge with the next newsletter.





European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

20th European Consensus Conference (EuCC) 2025

1 March 2025

Prepared by: Jörg Neugebauer, PhD, DMD

Chairman: Professor J. Neugebauer (Germany)

Protocol: Professor J. Neugebauer (Germany)

Participants: Ch. Berger (Germany)
Dr E. Cerekja (Albania)
Dr V. Gowd (India)
Dr A. Hartmann (Germany)
Dr F. Kasapi (Macedonia)
Professor P. Kobler (Croatia)
Professor V. Konstantinovic (Serbia)
Dr S. Liepe (Germany)
Dr W. Neumann (Germany)
Professor H. Özyuvaci (Turkey)
Dr G. Schioli (Italy)
Dr B. Singh (Nepal)
DDr M. Tröltzsch (Germany)
Dr J.W. Vaartjes (The Netherlands)
Professor J.E. Zöller (Germany)

Content

1 Methods	2
2 Problem	2
3 Patient sections	3
4 Surgical techniques	4
5 Prosthetic procedures	5
6 Conclusion	6
7 References	7



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

1 Methods

Objective

The purpose of this guide is to provide recommendations for clinicians active in implant dentistry, enabling the prevention, early detection and treatment of complications in order to optimize the patient outcome.

Introduction

All consensus recommendations in this paper should be interpreted as guidelines only. The patient's specific situation is always an important aspect and may justify deviations from the recommendations of this consensus paper.

Background

Implant placement is a proven way to replace missing teeth and to restore function and aesthetics. Nevertheless, complications may occur at various stages of treatment flow. Earlier guidelines covered surgical complications that might be harmful to anatomical structures; a risk analysis; and avoiding implant malpositioning considering further therapeutic needs. This guideline focuses on less frequently encountered risk factors that may arise at various treatment stages.

Literature search

The Cochrane Library, EMBASE, DIMDI and Medline literature databases were used to conduct a systematic search of recent published data. Selective search criteria were used, including terms such as

complication, dental implant, meta-analysis.

The publications identified with the search were screened by reading their abstracts and those irrelevant to the subject were identified and excluded. Those articles found to be potentially relevant were obtained in full-text form. Multiple review papers with meta-analyses were available on the subject.

Procedure for developing the guideline/consensus conference

A preliminary version of this document on which the EuCC based its deliberations was prepared and authored by Professor J. Neugebauer, University of Cologne, Faculty of Medicine and University Hospital Cologne, Department for Oral and Craniomaxillofacial and Plastic Surgery. The preliminary report was reviewed and discussed by the committee members in five steps, as follows:

- Reviewing the preliminary draft
- Collecting alternative proposals
- Voting on recommendations and levels of recommendation
- Discussing non-consensual issues
- Final voting

2 Problem

The outcome of implant therapy depends on the health status of the patient, including medication and nutritional status and the planned procedures and prosthetic restorations. From a surgical point of view, the use of surgical guides and grafting procedures may lead to complications. The recommendations for immediate loading require a high insertion torque, which is also a possible risk factor. From a restorative point of view, the retention type of the superstructure may be associated with technical or biological complications



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

or aesthetic failures [40]. The question of joining natural and implant abutments has also been subject to controversial discussion.

3 Patient sections

Patient expectations

The high number of implant treatments performed today may have deceived patients into believing that there are no longer any contraindications to implant treatment. Implants require a physiological bone metabolism, something that is not a given in the presence of several systemic diseases such as osteopetrosis (Morbus Albers-Schönberg), osteodystrophia deformans (Paget's disease of bone) or fibrous dysplasia. The bone metabolism can also be affected by medications, smoking habits or nutritional status.

Current observations

Case reports have stated that implant treatment is possible for patients with Paget's disease or fibrous dysplasia. For patients receiving antiresorptive therapy, a high incidence of complications in the form of bone necrosis has been reported after tooth extraction, surgical interventions or even as a result of sore spots. However, implant placement, possibly in conjunction with autologous grafting procedures, could produce positive outcomes in osteoporosis patients [32]. There is a need to develop specific treatment strategies for each individual patient [28].

Low level of cholecalciferol (vitamin D3) may compromise osseointegration and graft regeneration or lead to progressing peri-implantitis [6]. Patients receiving proton pump inhibitors (PPI) or serotonin reuptake inhibitors (SRI) exhibit higher rates of implant failure [16]. Conflicting results have been reported regarding the effect of glucocorticoids and NSAIDs on implant treatment outcomes [12].

Preventions of complications

- Implant placement is contraindicated in patients suffering from osteopetrosis.
- High-dosage antiresorptive therapy could result in higher rate of BRONJ [35].
- Patients who have been on antiresorptive therapy for osteoporosis need a detailed case selection with surgical techniques not requiring intensive bone remodelling [15, 35]. Extensive bone splitting, osteotome techniques or lateral sinus grafts should be avoided.
- No increased general risk for patients suffering from osteoporosis has been reported [20].
- In patients with soft bone evident in preoperative radiographs or with increased bone resorption, blood cholecalciferol levels should be checked [6].
- In patients with PPI or SRI, the duration and amount of drugs could be investigated before considering a patient for implant treatment [16].
- Patients must be informed that untreated periodontitis, uncontrolled diabetes or smoking may substantially increase the risk for biologic complications (e.g. peri-implantitis) [3, 7, 23, 33].
- In patients under long-term glucocorticoid medication at high doses, bone-metabolism parameters may have to be evaluated.
- Mild bone malformation in patients with fibrous dysplasia or Paget's disease need a strict indication for dental implants due to a lack of pertinent data.
- Depending on the surgical indication, patient may profit by an antibiotic prophylaxis for reduced (early) implant failure [13, 36, 38].



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

4 Surgical techniques

Patient expectations

Patients increasingly request immediate fixed rehabilitation in conjunction with immediate implant placement and loading. However, postoperative morbidity should be kept as low as possible.

Current observations

High insertion torque

Immediate implant placement with immediate restoration is a scientifically proven treatment concept for rehabilitating a failing dentition [10]. Various recommendation on the determination of primary stability have been given, depending on implant designs and the surgical procedures performed, for achieving osseointegration in the context of immediate restoration [37]. A recent RCT on insertion torque showed increased failure and bone resorption rates in the mandible for high insertion torques [25]. Previous meta-analyses have shown that high insertion torques are not correlated with increased bone resorption or implant failure [4, 22].

Flapless surgery

Implant placement using 3D surgical guides is now established, and flapless surgery should reduce the postoperative discomfort. The use of surgical guides based on CBCT technology permits highly accurate implant placement [5, 8]. Compared to free-hand flapless surgery and to the raising of a flap, the outcome of guided flapless surgery was not different in terms of implant failure rates and bone resorption in the hands of experienced treatment providers [21, 41]. Nevertheless, complication such as bone perforation or displacement of the surgical guide may occur [5, 8, 30].

Soft-tissue grafting

To achieve better outcomes in the aesthetic zone, soft-tissue grafting should be considered, particularly for immediate implant placement [31]. Grafting may also be necessary in case of soft-tissue dehiscence [2].

Amount of grafting

Grafting procedures carry a certain risk of complications. The use of shorter implants could reduce the amount of grafting required, thereby potentially lowering these risks [1].

Prevention of complications

- Due to the many different implant designs and recommend preparation techniques, especially in dense bone or in the presence of a thin cortical plate, the manufacturers' recommend insertion torques should be considered.
- Patients benefit from flapless procedures if a proper indication exists in terms of the available bone supply and preoperative 3D diagnostic findings.
- Flapless procedures are subject to a specific learning curve.
- Soft-tissue grafting reduces the risk of aesthetic complications.



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

5 Prosthetic procedures

Patient expectations

Patients expect a stable prosthetic restoration that meets their aesthetic and functional needs, with minimal complications.

Current observations

A reduction in the number of implants in a given case due to economic or anatomical reasons may be considered by using both teeth and implants as abutments for fixed partial dentures (FPDs). Superstructures can be cemented or screw-retained, both of which can be associated with complications. In recent years a large spread of materials, prosthetic designs and components are used.

Tooth-to-implant fixed partial dentures

A meta-analysis of tooth-to-implant (hybrid) fixed partial dentures (T-I FPDs) reported survival rates of 94.1% after 5 years and 77.8% after 10 years of clinical service [18]. The impact of T-I FPDs and implant-to-implant FPDs in the partially edentulous arch on implant survival rates showed no significant differences for periods up to 72 months [27, 42]. A recent systematic review assessed the effect of rigid and non-rigid splinting between implants and teeth, with overall prosthetic survival rates of 85% and higher risks for tooth intrusion associated with non-rigid connections for observation periods of between 18 and 120 months [39].

Retention of superstructures

Depending on the number of implants and the design of available abutments, superstructures can be cemented or screw-retained. Technical or biological complications may occur with either type of retention. A meta-analysis showed no differences regarding loosening of superstructures, changes in marginal bone levels or peri-implantitis [19, 24]. Using 6 implants for full-arch restoration in the maxilla result in fewer complications compared to using 4 implants [34].

Design of superstructures

To minimize the number of implants, cantilevers are often used in prosthetic reconstructions. However, these superstructures are associated with a higher incidence of technical complications, particularly in full-arch reconstructions [17] and single crowns [9, 17]. Patients suffering from bruxism have also been found to experience more frequent technical complications [14].

To reduce the risk of complications such as chipping, monolithic zirconia superstructures perform better than veneered zirconia in full-arch reconstructions [26, 29]. Additionally, metal-ceramic superstructures exhibit fewer complications compared to metal-acrylic resin implant-supported fixed complete dental prostheses [11].

Prevention of complications

- Rigid superstructures should be preferred for T-I FPDs.
- Complications associated with T-I FPDs are encountered mainly at the natural abutment, especially when the teeth are periodontally compromised or root canal filled.
- The form of retention of the superstructure should be chosen by taking function, aesthetics and professional maintenance into account rather than focusing on available techniques.
- To facilitate maintenance, a retrievable superstructure is preferred, but a definitive cementing on natural tooth.
- To avoid technical complication the biomechanical considerations for each patients must be evaluated.



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications in implant treatment

6 Conclusion

Dental implants are reliable treatment options for restoring patient function and aesthetics. Careful case selection is necessary by considering not only the oral findings alone. Due to the great variability of implant designs and surgical and prosthetic procedures proposed, the individual suggested parameter should be followed to avoid complication. All procedures should be performed by treatment providers with the requisite up-to-date expertise and training.

Cologne, 1 March 2025



Professor Joachim E. Zoller
Vice-President



Professor Jörg Neugebauer
Chairman of EuCC



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications to optimize patient

7 References

1. Abayov P, Sarikov R, Nazarenko LM et al. Outcome Difference between Short and Longer Dental Implants Placed Simultaneously with Alveolar Bone Augmentation: a Systematic Review and Meta-Analysis. *J Oral Maxillofac Res* 2024; 15: e2.
2. Amid R, Moscowchi A, Azizi S et al. Coverage stability of peri-implant soft tissue dehiscence: A systematic review and meta-analysis. *J Esthet Restor Dent* 2024; 36: 284-294.
3. Bencze B, Cavalcante BGN, Romandini M et al. Prediabetes and poorly controlled type-2 diabetes as risk indicators for peri-implant diseases: A systematic review and meta-analysis. *J Dent* 2024; 146: 105094.
4. Berardini M, Trisi P, Sinjari B et al. The Effects of High Insertion Torque Versus Low Insertion Torque on Marginal Bone Resorption and Implant Failure Rates: A Systematic Review With Meta-Analyses. *Implant Dent* 2016; 25: 532-540.
5. Bover-Ramos F, Vina-Almunia J, Cervera-Ballester J et al. Accuracy of Implant Placement with Computer-Guided Surgery: A Systematic Review and Meta-Analysis Comparing Cadaver, Clinical, and In Vitro Studies. *Int J Oral Maxillofac Implants* 2018; 33: 101-115.
6. Choukroun J, Khoury G, Khoury F et al. Two neglected biologic risk factors in bone grafting and implantology: high low-density lipoprotein cholesterol and low serum vitamin D. *J Oral Implantol* 2014; 40: 110-114.
7. Chrcanovic BR, Albrektsson T, Wennerberg A. Smoking and dental implants: A systematic review and meta-analysis. *J Dent* 2015; 43: 487-498.
8. Colombo M, Mangano C, Mijiritsky E et al. Clinical applications and effectiveness of guided implant surgery: a critical review based on randomized controlled trials. *BMC Oral Health* 2017; 17: 150.
9. De Angelis P, Manicone PF, Liguori MG et al. Clinical and radiographic evaluation of implant-supported single-unit crowns with cantilever extensions: A systematic review and meta-analysis. *J Prosthodont* 2024; 33: 841-851.
10. Del Fabbro M, Ceresoli V, Taschieri S et al. Immediate loading of postextraction implants in the esthetic area: systematic review of the literature. *Clin Implant Dent Relat Res* 2015; 17: 52-70.
11. Estrin N, Nam K, Romanos GE et al. Clinical Outcomes of Metal-Ceramic versus Metal-Acrylic Resin Implant-Supported Fixed Complete Dental Prostheses: A Systematic Review and Meta-analysis. *Int J Prosthodont* 2023; 36: 354-365.
12. Fu JH, Bashutski JD, Al-Hezaimi K et al. Statins, glucocorticoids, and nonsteroidal anti-inflammatory drugs: their influence on implant healing. *Implant Dent* 2012; 21: 362-367.
13. Ikram J, Shado R, Pereira IN et al. Efficacy and safety of using antibiotics to prevent post-operative complications in oral implant treatment: evidence-based review. *BDJ Open* 2023; 9: 47.
14. Ionfrida JA, Stiller HL, Kammerer PW et al. Dental Implant Failure Risk in Patients with Bruxism-A Systematic Review and Meta-Analysis of the Literature. *Dent J (Basel)* 2024; 13.
15. Jung J, Shim GJ, Park JS et al. Effect of anti-resorptive therapy on implant failure: a systematic review and meta-analysis. *J Periodontal Implant Sci* 2024.
16. Jung RE, Al-Nawas B, Araujo M et al. Group 1 ITI Consensus Report: The influence of implant length and design and medications on clinical and patient-reported outcomes. *Clin Oral Implants Res* 2018; 29 Suppl 16: 69-77.
17. Kondo Y, Sakai K, Minakuchi H et al. Implant-supported fixed prostheses with cantilever: a systematic review and meta-analysis. *Int J Implant Dent* 2024; 10: 57.
18. Lang NP, Pjetursson BE, Tan K et al. A systematic review of the survival and complication rates of fixed partial dentures (FPDs) after an observation period of at least 5 years. II. Combined tooth-implant-supported FPDs. *Clinical oral implants research* 2004; 15: 643-653.
19. Lemos CA, de Souza Batista VE, Almeida DA et al. Evaluation of cement-retained versus screw-retained implant-supported restorations for marginal bone loss: A systematic review and meta-analysis. *J Prosthet Dent* 2016; 115: 419-427.
20. Lemos CAA, de Oliveira AS, Fae DS et al. Do dental implants placed in patients with osteoporosis have higher risks of failure and marginal bone loss compared to those in healthy patients? A systematic review with meta-analysis. *Clin Oral Investig* 2023; 27: 2483-2493.
21. Lemos CAA, Verri FR, Cruz RS et al. Comparison between flapless and open-flap implant placement: a systematic review and meta-analysis. *Int J Oral Maxillofac Surg* 2018.
22. Li H, Liang Y, Zheng Q. Meta-Analysis of Correlations Between Marginal Bone Resorption and High Insertion Torque of Dental Implants. *Int J Oral Maxillofac Implants* 2015; 30: 767-772.
23. Li Y, Lu Z, Sun H. Impact of diabetes mellitus on the poor prognosis in patients with osseointegrated dental implants: a meta-analysis of observational studies. *Biotechnol Genet Eng Rev* 2024; 40: 473-491.
24. Ma S, Fenton A. Screw- versus cement-retained implant prostheses: a systematic review of prosthodontic maintenance and complications. *Int J Prosthodont* 2015; 28: 127-145.
25. Marconcini S, Giammarinaro E, Toti P et al. Longitudinal analysis on the effect of insertion torque on delayed single implants: A 3-year randomized clinical study. *Clin Implant Dent Relat Res* 2018; 20: 322-332.



European Association of Dental Implantologists

Bundesverband der implantologisch
tätigen Zahnärzte in Europa e.V.

Guidelines 2025

Update: Preventing, detecting and treating specific complications to optimize patient

26. Martakoush-Saleh S, Salgado-Peralvo AO, Pena-Cardelles JF et al. Evaluating the clinical behavior of veneered zirconia in comparison with monolithic zirconia complete arch implant-supported prostheses: A systematic review and meta-analysis. *J Prosthet Dent* 2024; 132: 899-912.
27. Muddugangadhar BC, Amarnath GS, Sonika R et al. Meta-analysis of Failure and Survival Rate of Implant-supported Single Crowns, Fixed Partial Denture, and Implant Tooth-supported Prostheses. *J Int Oral Health* 2015; 7: 11-17.
28. Penoni DC, Canellas JVS, Silami M et al. Osteonecrosis of the jaws in patients under osteoporosis treatment: a nine-year experience report. *Arch Endocrinol Metab* 2023; 67: e000612.
29. Pjetursson BE, Sailer I, Merino-Higuera E et al. Systematic review evaluating the influence of the prosthetic material and prosthetic design on the clinical outcomes of implant-supported multi-unit fixed dental prosthesis in the posterior area. *Clin Oral Implants Res* 2023; 34 Suppl 26: 86-103.
30. Romandini M, Ruales-Carrera E, Sadilina S et al. Minimal invasiveness at dental implant placement: A systematic review with meta-analyses on flapless fully guided surgery. *Periodontol* 2000 2023; 91: 89-112.
31. Rondone EM, Leitao-Almeida B, Pereira MS et al. The Use of Tissue Grafts Associated with Immediate Implant Placement to Achieve Better Peri-Implant Stability and Efficacy: A Systematic Review and Meta-Analysis. *J Clin Med* 2024; 13.
32. Schmitt CM, Buchbender M, Lutz R et al. Oral implant survival in patients with bisphosphonate (BP)/antiresorptive and radiation therapy and their impact on osteonecrosis of the jaws. A systematic review. *Eur J Oral Implantol* 2018; 11 Suppl 1: S93-S111.
33. Serroni M, Borgnakke WS, Romano L et al. History of periodontitis as a risk factor for implant failure and incidence of peri-implantitis: A systematic review, meta-analysis, and trial sequential analysis of prospective cohort studies. *Clin Implant Dent Relat Res* 2024; 26: 482-508.
34. Sharaf MA, Wang S, Mashrah MA et al. Outcomes that may affect implant and prosthesis survival and complications in maxillary fixed prosthesis supported by four or six implants: A systematic review and meta-analysis. *Heliyon* 2024; 10: e24365.
35. Stavropoulos A, Bertl K, Pietschmann P et al. The effect of antiresorptive drugs on implant therapy: Systematic review and meta-analysis. *Clin Oral Implants Res* 2018; 29 Suppl 18: 54-92.
36. Tan LW, Ng YE, Giok KC et al. Comparative Efficacy of Different Amoxicillin Dosing Regimens in Preventing Early Implant Failure-A Systematic Review with Network Meta-Analysis. *Antibiotics (Basel)* 2023; 12.
37. Tettamanti L, Andrisani C, Bassi MA et al. Immediate loading implants: review of the critical aspects. *Oral Implantol (Rome)* 2017; 10: 129-139.
38. Torof E, Morrissey H, Ball PA. Antibiotic Use in Dental Implant Procedures: A Systematic Review and Meta-Analysis. *Medicina (Kaunas)* 2023; 59.
39. Tsaousoglou P, Michalakakis K, Kang K et al. The effect of rigid and non-rigid connections between implants and teeth on biological and technical complications: a systematic review and a meta-analysis. *Clin Oral Implants Res* 2017; 28: 849-863.
40. Verma A, Singh SV, Arya D et al. Mechanical failures of dental implants and supported prostheses: A systematic review. *J Oral Biol Craniofac Res* 2023; 13: 306-314.
41. Voulgarakis A, Strub JR, Att W. Outcomes of implants placed with three different flapless surgical procedures: a systematic review. *Int J Oral Maxillofac Surg* 2014; 43: 476-486.
42. Weber HP, Sukotjo C. Does the type of implant prosthesis affect outcomes in the partially edentulous patient? *Int J Oral Maxillofac Implants* 2007; 22 Suppl: 140-172.

THE KING

AMONG THE BEGO IMPLANTS

BEGO Semados® RSX^{Pro}



MAKE THE MOVE OF YOUR LIFE!

The BEGO Semados® RSX^{Pro} implant is more than just an implant. It is a guarantee of quality, reliability, and first-class results. With its versatility and adaptability, it is the perfect solution for a wide range of dental applications. Discover the future of dental implantology!



Curious?

<https://www.bego.com/king>

