

Immediate anterior maxilla restoration

Integrating guided bone regeneration and digital planning for aesthetic results

Dr Dalton Marques, Brazil

The following case report focuses on the rehabilitation of a 35-year-old male patient who presented with loose anterior teeth, a progressively widening diastema, and discomfort during eating. The patient desired a comprehensive solution to restore dental stability, improve oral function, and enhance his smile. His dental history included prior orthodontic treatment for an impacted canine, with no systemic conditions, smoking habits, or medication use influencing the treatment plan.

Restoration of the aesthetic zone in a young patient is particularly challenging due to the high expectations for natural-looking results and the functional demands placed on the anterior teeth. In addition to aesthetic considerations, these cases often involve compromised bone and soft-tissue conditions, requiring precise surgical and prosthetic planning. Immediate treatment options in such scenarios provide a clear advantage by reducing treatment time, preserving soft-tissue contours, and main-

taining patient confidence during the rehabilitation process.

The use of Straumann® BLX implants was vital in addressing the complexity of this case. These implants offer optimal primary stability, even in situations with limited bone availability. Their design facilitates efficient osseointegration while supporting soft-tissue health, making them a reliable choice for restoring the aesthetic zone. The immediate placement approach minimised treatment duration

and preserved the natural gingival architecture, contributing to an optimal aesthetic outcome.

Guided bone regeneration (GBR) played a crucial role in the treatment due to the significant vertical and horizontal bone loss identified through radiographic assessment. cerabone®, a deproteinised bovine bone substitute, was employed to fill the gaps and support the osseointegration of the implants. Known for its excellent biocompatibility and structural stability, cerabone® helps preserve ridge volume and ensures long-term support for soft and hard tissues. Its slow resorption rate is particularly advantageous in the aesthetic zone, where maintaining ridge contours is essential for achieving natural-looking results.

Following a thorough clinical evaluation and classification of the case as complex for surgery and advanced for prosthodontics under SAC criteria, a digital workflow was integrated into the treatment plan. This approach allowed for precise planning of implant placement and prosthetic design, ensuring a functional and aesthetic restoration.

This report highlights the challenges of managing the aesthetic zone in a young patient and the advantages of combining



Figs. 1a+b: Hopeless maxillary central and lateral incisors.

immediate implant placement, biomaterials, and digital technology. The integration of these elements highlights the potential to achieve stable, predictable, and natural-looking outcomes that meet both functional demands and the patient's expectations.

Initial situation

A 35-year-old male patient presented with the chief complaint of loose front teeth and pain, especially while eating. He expressed the desire to achieve a stable dentition, improve function, and have a more aesthetic smile. His dental history includes orthodontic treatment performed years ago due to an impacted canine. There were no reports of systemic diseases, smoking, or medication use.

The patient presented a low smile line with no signs of local inflammation. There was mobility grade II in the front teeth and a diastema centralis, which, according to the patient, has progressively widened over time (Fig. 1).

The radiographic evaluation showed loss of vertical and horizontal bone, as well as a thin buccal plate (Figs. 2–4).





Based on the SAC classification, the patient's surgical case was categorised as complex, while the prosthodontic status was classified as advanced (Fig. 5).

The prognosis of the remaining teeth, based on clinical and radiographic analysis, was poor. Extraction of hopeless teeth was necessary as part of the treatment plan.

Treatment planning

After discussing the treatment alternatives with the patient, including the advantages, disadvantages, risks, and complications, the decision was made to opt for:

1. Digital planning of a fixed immediate rehabilitation on three implants.
2. Extraction of hopeless teeth #13, #12, #11, #21 and #22.
3. Immediate placement of three Straumann® BLX Roxolid® SLActive® implants in positions #13 (5.0 x 10 mm), #11 (3.75 x 12 mm), and #22 (4.5 x 10 mm).



Figs. 2–4: Radiograph showing vertical and horizontal bone loss with a thin buccal plate. – **Fig. 5:** SAC classification.

4. Bone grafting with cerabone® deproteinised bovine bone substitute.
5. Delivery of screw-retained temporary prosthesis.
6. Delivery of screw-retained definitive prosthesis.

Surgical procedure

Local anesthesia with lidocaine 2% with epinephrine 1:100,000 was administered. The hopeless teeth #13, #12, #11, #21, and #22 were extracted atraumatically. A mucoperiosteal flap with a crestal inci-

sion was made, and a surgical stent was positioned to guide implant-site preparation following the extractions. The implant beds were prepared using the Straumann® BLX Surgical Cassette following the manufacturer's instructions (Fig. 6).

Immediate BLX Straumann® implants were placed in positions #13 (5.0 x 10 mm), #11 (3.75 x 10 mm), and #22 (4.5 x 10 mm; Fig. 7). The gaps and sockets were carefully filled with cerabone® deproteinised bovine bone substitute, small granules, to improve osseointegration and provide permanent structural support (Fig. 8).






Fig. 6: Implant bed preparation following extraction of hopeless teeth. – **Fig. 7:** Immediate Straumann® BLX implants placed at #13 (5.0 × 10 mm), #11 (3.75 × 10 mm), and #22 (4.5 × 10 mm). – **Fig. 8:** Gaps and sockets filled with cerabone® granules. – **Fig. 9:** The flap is closed with sutures.

This bone substitute was selected for its excellent biocompatibility and its capacity to support soft tissue in the aesthetic area, helping to preserve the shape of the ridge.

Sutures were then placed to close the flap, promoting optimal healing conditions (Fig. 9).

Prosthetic procedure

Digital impressions were taken using the Straumann® Virtuo Vivo™ intra-oral scanner and a five-unit bridge was digitally planned (Fig. 10). Additionally, an occlusal view of the virtually planned five-unit bridge was generated to ensure precise alignment and aesthetics (Fig. 11).

A temporary bridge was then fabricated using a milled resinous material on the Straumann® CARES® C series milling machine (Fig. 12). The screw-retained temporary bridge was then delivered to the

patient, providing functional and aesthetic restoration while supporting tissue healing and adaptation (Fig. 13).

At the six-month follow-up, healing was uneventful. The final restoration was also digitally planned (Fig. 14). The final screw-retained bridge was delivered, occlusion was checked, and instructions were given. The patient was very satisfied with the results (Fig. 15).

Treatment outcomes

Eighteen months after delivery of the definitive prosthesis, clinical and radiographic evaluation demonstrated stable implants, healthy peri-implant bone, and satisfactory prosthesis integration (Figs. 16+17). At the three-year follow-up, these results remained stable, with no complications observed (Figs. 18–21).




Fig. 10: Digital impressions with Straumann® Virtuo Vivo™ and five-unit bridge digitally planned. – **Fig. 11:** Occlusal view of digitally planned five-unit bridge.



Fig. 12: Temporary bridge fabricated on the Straumann® CARES® C series. – **Fig. 13:** Screw-retained temporary bridge delivered. – **Fig. 14:** Digitally planned final restoration. – **Fig. 15:** Final screw-retained bridge delivered. – **Figs. 16+17:** Eighteen months follow up: clinical and radiographic evaluation showing satisfactory functional and aesthetic outcome. – **Figs. 18–21:** Three-year follow-up: stable clinical and radiographic results without complications.

The patient expressed high satisfaction, noting restored function, confidence in smiling and speaking, and a significant improvement in quality of life, highlighting the value of dental implant therapy in comprehensive oral rehabilitation.

Conclusion

Rehabilitation in the anterior area of the maxilla becomes a challenge due to

the aesthetic and functional requirements. In cases with a complex aesthetic and functional deficiency, proper planning is a priority. The precision and efficiency of the digital workflow enable accurate prosthetic design, implant position, and determine whether additional surgical procedures are required.



Corresponding author

Dr Dalton Marques

Curitiba, Brazil

+55 41 991993601

dalton_marques@hotmail.com

www.drdaltonmarques.com.br