

Practical periodontics in daily practice

Author_Dr Amit Patel, UK



Fig. 1 Inflamed gingival tissue and pocketing around a bridge abutment.



Fig. 2



Fig. 3



Fig. 4

Fig. 1 Inflamed gingival tissue and pocketing around a bridge abutment.

Fig. 2 Healthy-looking gingival tissues, but the patient is a smoker and 5 mm pocketing is noted.

Fig. 3 BPE probe.

Fig. 4 Williams' probe.

_Cosmetic and implant dentistry has become increasing popular among dentists and patients. As this type of dentistry increases, so does litigation. Legal defence organisations have noted that their highest litigation costs are due to an increase in undiagnosed periodontal disease, as well as poorly planned cosmetic and implant dentistry. A failure to diagnose periodontal disease, inadequate records, poor quality treatment and treatment planning, supervised neglect, and failure to refer all lead to increased litigation within the profession.

The guidance for standards set by the General Dental Council states that a clinician should work within his or her knowledge, professional competence and physical abilities, should refer patients for a second opinion and for further advice when necessary, and should refer patients for further treatment when necessary.

of the periodontium is stable and good plaque control has been achieved, the restorative treatment will have a better long-term prognosis. This article will briefly discuss the simple tools we have in our surgeries to help diagnose periodontal disease and when to treat and when to refer using the British Society of Periodontology's referral guidance.

The clinical signs of chronic periodontal disease are gingival inflammation and bleeding, pocketing, gingival recession, tooth mobility and migration, alveolar bone loss and halitosis. Figure 1 shows a patient with gingival inflammation and bleeding on probing with a pocket depth of greater than 5 mm. A good predictor of gingival health is no bleeding on probing, but it is important to note that in smokers the gingival tissues look relatively healthy and in most cases do not bleed on probing, as smoking masks the presence of disease (Fig. 2).

An essential tool for the assessment of the periodontal tissues is the BPE (basic periodontal examination) probe, otherwise known as a WHO 621 probe (Fig. 3). This probe has a ball tip of 0.5 mm in diameter

Table I _ Basic periodontal examination codes.

Fig. 5 Furcation probe.

Fig. 6 Long cone radiograph of teeth 14 to 17.

It is important as the clinician to assess the periodontal condition before starting any restorative dentistry, whether simple or complex. There is very good long-term evidence that once the foundation

Code Signs	Treatment
0 No bleeding on probing Pocket depth of ≤ 3 mm	No treatment
1 Bleeding on probing	Hygiene instruction
2 Plaque retentive factors present No pocket depth of > 3.5 mm	Hygiene instruction and scaling
3 Pocket depth of > 3.5 mm but < 5.5 mm	Hygiene instruction, supra- and subgingival scaling
4 Pocket depth of ≥ 5.5 mm	Full periodontal assessment
* Furcation	Full periodontal assessment





Fig. 5



Fig. 6



and a black band marked at 3.5–5.5 mm. The BPE is a simple and quick way of screening our patients for any underlying periodontal disease. Williams' probes are also commonly used to assess the periodontal tissue (this probe has graduated markings at 1–2, 3–5, 7–8, 9–10 mm; Fig. 4).

The BPE was developed by the British Society of Periodontology. It is a method of screening patients to determine the treatment required for the level of disease present. The BPE is an index for treatment need and does not estimate the level of disease present. Rather, a loss of attachment chart is used to determine this.

The BPE divides the mouth into sextants. All the teeth in a sextant are examined and scored accordingly (Table I). Recently, the British Society of Periodontology has made a slight change to the BPE scoring. Code* has now been changed to denote the presence of a furcation only, whereas previously it was to denote the presence of a furcation or attachment loss of 7 mm or greater. The society has also said that the other BPE codes and Code* should be recorded for each sextant where a furcation is present. An example is shown in Table II.

A furcation probe or Nabers' probe is also an essential tool for assessing the degree of furcation involvement of a molar tooth. We can measure the amount of horizontal bone loss that has occurred within the furcation, classifying it as a Class 1, 2 or 3 furcation. The dark bands represent 3 mm markings (Fig. 5). A Class 1 furcation is noted when the furcation probe penetrates less than 3 mm into the furcation (Fig. 6). A Class 2 furcation is when the probe penetrates greater than 3 mm but does not go all the way through the furcation

(Figs. 7 & 8). A Class 3 furcation is when the probe passes through the furcation unimpeded (Figs. 9 & 10). Radiographs are another important tool used to assess the bone levels around each tooth, root morphology and furcation involvement, and therefore the support present and long-term prognosis of the teeth (Fig. 8). Long-cone parallel radiographs or vertical bitewings are taken of sextants when the score is 3 or more (Fig. 11).

_Risk factors

The clinician should also be aware of risk factors that can exacerbate the existing periodontal disease: diabetes, smoking and genetics. A combination of these factors makes certain patients susceptible to higher risks of periodontal disease. These cases may be treated in practice but referral to a specialist would be required if the disease is not stabilised. Recognition of risk factors includes:

- _BPE Codes 3, 4 or * in patients under 35 years old;
- _smoking ten or more cigarettes a day;
- _a medical condition directly affecting the periodontal tissues, for example diabetes, stress and certain types of medication;
- _a root morphology that adversely affects prognosis;
- _rapid periodontal breakdown (> 2 mm attachment loss in any one year);
- _a high bleeding percentage with a low plaque score; and
- _a family history of early tooth loss due to periodontal disease.

_Oral hygiene

A high standard of oral hygiene is critical for successful periodontal therapy. There is a great deal

Fig. 7_ Class 1 buccal furcation in tooth 16.
Fig. 8_ Class 2 buccal furcation in tooth 17.
Fig. 9_ Class 2 lingual furcation in tooth 47.
Fig. 10_ Class 3 furcation in tooth 46.
Fig. 11_ Class 3 furcation in tooth 46.

4	2	4*
3*	3	2

Table II_ Example of BPE scoring and Code* for each sextant where a furcation is present.




Fig. 12_ Interdental brushes.
Fig. 13_ Interdental brush used to maintain a Class 2 buccal furcation.
Fig. 14_ Consultation visit.
Fig. 15_ Long-cone periapical radiograph of teeth #43 to 46.



Fig. 16_Tooth #46 with 10 mm pocketing mesially.

Fig. 17_Tooth #45 with 9 mm pocketing mesially.

Fig. 18_Largest interdenal brush for tooth #46 demonstrated to patient.

Fig. 19_Largest interdenal brush for tooth #45.

Fig. 20_Tooth #46 with 6 mm pocketing mesially after simple oral hygiene instruction given (large interdenal brushes).

Fig. 21_Tooth #45 with 7 mm pocketing mesially after simple oral hygiene instruction given (large interdenal brushes).

Fig. 22_Teeth #43 to 46 show increased recession and a reduction in probing depth to 3 mm after non-surgical therapy.

Fig. 23_Gingival health after initial therapy with a reduction in probing depth.

of evidence that regular plaque removal around periodontally involved teeth at a level that prevents bleeding on probing leads to a reduction in disease progression. It is essential that the patient be taught simple, yet effective ways to improve his or her plaque control at home with daily use of a rotating-oscillating electric toothbrush and interdenal brushes (Fig. 12). A systematic review by the Cochrane Library has shown that a rotating-oscillating electric toothbrush is far more effective at removing plaque.¹

A study has shown that using the correct size interdenal brushes can improve the periodontal condition significantly.² A recent systematic review has also shown that flossing by patients has no effect on the plaque index or gingival index and that flossing is not effective in periodontally compromised patients. Interdenal brushes have been shown to remove more plaque than flossing.³

It is important to motivate your patients to use the largest interdenal brushes (Figs. 12 & 13), but this can be a little difficult because they may not be able to see the short-term benefits, as their gums may bleed more and the interproximal spaces will become larger. It is essential to reinforce the same message. This will reassure the patient and after a short time they will see visible benefits, that is, less bleeding on cleaning and a healthier-looking gingiva.

Case report

A 32-year-old male patient was referred to me complaining of loose teeth and bleeding gums for over 12 months. He was fit, apparently healthy and a non-smoker. A diagnosis of generalised aggressive periodontitis was made from the clinical examination (Figs. 14–17). At the consultation appointment, oral hygiene instruction was given and the largest interdenal brushes demonstrated. At the following appointment, the full-mouth non-surgical phase was carried out with systemic antibiotics.⁴ From Figures 18 to 21, it can be noted that using the

correct size interdenal brushes can lead to a reduction in inflammation and therefore a reduction in pocket depth.

Eight weeks after the non-surgical phase, the patient was reviewed for a periodontal reassessment. It was noted that the periodontal tissue had responded extremely well to the initial therapy with pocket depth of 3 to 4 mm throughout the mouth (Figs. 22 & 23). At this point, the patient would be placed on a three-monthly maintenance regime, with reinforcement of oral hygiene instruction and subgingival plaque removal for any deep sites. This would be carried out by his general dentist or hygienist. My plan would be to review the patient in six months' time to review his periodontal condition.

It is important to know that periodontal therapy works and a healthy periodontium is the backbone of good restorative dentistry. Treating periodontal disease can be challenging but can also be very rewarding. Careful assessment, treatment, referral to a specialist if necessary and monitoring of your patients are essential for avoiding any future problems.

For further information regarding the BPE and referral guidance, contact the British Society of Periodontology or refer to its website: www.bsperio.org.uk.

Editorial note: A complete list of references is available from the publisher.

about the author

cosmetic
dentistry



Dr Amit Patel
Specialist in Periodontics & Implant Dentist, Associate Specialist Birmingham Dental Hospital, Honorary Clinical Lecturer University of Birmingham School of Dentistry



Tribune CME

Clinical Masters Program in Aesthetic and Restorative Dentistry

10-14 January 2013 and 24-27 April 2013 in Dubai, for a total 9 days



Didier Dietschi



Francesco Mangani



Panos Bazos

Session I: 10 - 14 January 2013 (5 days)

- Direct/Indirect composite Artistry in the Anterior Segment
- Direct/Indirect composite Artistry in the Posterior Segment
- Photography and shade analysis

Clinical Masters:

Didier Dietschi, Francesco Mangani, Panos Bazos



Mauro Fradeani



Urs Brodbeck

Session II: 24 - 27 April 2013 (4 days)

- Full coverage Anterior/Posterior Restoration
- Partial coverage Anterior/Posterior Restoration, Ceramic Restoration

Clinical Masters:

Mauro Fradeani, Urs Brodbeck

ADA C.E.R.P.® Continuing Education
Recognition Program

Tribune America LLC is the ADA CERP provider.
ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry.

Dubai, UAE

2 on location sessions with **live patient treatment**
and **hands on** in each session!
+ online live and recorded lectures, **online mentoring**



+ Full access to our online learning platform:
hours of premium video training and live webinars
collaborate with peers and faculty on **your cases**

Registration info

Course fee: € 8,900

Register by the end of October and you
will receive a complimentary iPad
preloaded with premium dental courses

Tribune CME

Tel.: +49-341-48474-302

email: info@tribunecme.com

www.TribuneCME.com



Discover the Master's secrets and Dubai's superlatives

