

Lateral maxillary incisor implant—Key issues for aesthetic success

Authors: Drs Philippe Russe & Patrick Limbour, France



Fig. 1

Fig. 2

Introduction

Fig. 1: Agenesis of 22, opening of orthodontic space.

Fig. 2: Line of intermediate smile. The smile uncovers the papillae and reaches the collar of the incisors (12 and 22 are supported by implants).

Faced by a missing lateral incisor, practitioners often consider a wide range of issues and are also faced by numerous treatment options:

- in a young patient, faced with a unilateral or bilateral agenesis, he has to choose between an orthodontic treatment that either opens up the spaces or closes them. This decision, when taken early in the overall treatment, will affect both the patient and their caregiver for a long time (Fig. 1);

- in an adult patient, this is a consequence of bone, physiological, traumatic or infectious resorption, which will result in a decision whether or not to recommend a bone reconstruction or a gingival augmentation.

In every situation, the results will be judged by the patient and those around him. Since the lateral maxillary incisor is an integral part of the smile, aesthetic expectations are generally very high and, if the results do not meet the expectations, disappointment can be powerfully felt.

Fig. 3a: Average forms, types and dimensions of the lateral incisor according to Papathanassiou.⁶

Overall height: 21 mm, coronal height: 9 mm, radical height: 12 mm, mesiodistal cervical diameter: 5 mm, mesiodistal coronal diameter: 6.5 mm, vestibular-lingual cervical diameter: 5 mm, vestibular-lingual coronal diameter: 6.5 mm.

Fig. 3b: Proximal view photographs showing ten anatomical variants of lateral maxillary incisors described by the author.⁶




Fig. 3a




Fig. 3b



Fig. 4




Fig. 5

When describing the different treatment stages, a number of pitfalls and difficulties will be highlighted and advice and clinical protocols will be given, in order to ensure that the results of this implant/prosthetic treatment are predictable and as aesthetically attractive as possible. This first article is concerned with these issues as regards the pre-prosthetic stages; the second will consider the most important aspects of the prosthetic stages as well as aesthetic outcomes and their evolution over the long term.

Anamnesis

Once the usual contraindications for oral and implant surgery have been eliminated, particular attention should be given to the patient's answers concerning their smoking habits. Indeed, meta-analysis give an accurate picture of the consequences of smoking, with increases of:

- peri-implantitis^{1,2} and bone loss²;
- failure rates.³

The conclusions of Snider et al.⁴ can provide recommendations for the practitioner faced with a patient who is a smoker:

- the best is to ask the patient to stop smoking...;
- if this approach is not acted on, then the patient must be warned of the increased risk of failure and of postoperative complications.

This last issue is important, as smoking can be considered a lost opportunity as far as implant treatment is concerned.

“It is preferable to avoid patients that are smokers.”

Clinical examination

The smile line

When replacing a tooth in an aesthetic region, understanding the location of the smile line is one of the determining issues during the clinical examination. There are two factors to consider: the exposure of papillae and visibility of the collar of the lateral incisor, and there is one significant problem: any aesthetic deficit experienced by the patient tends to make them change their smile line, which can happen more or less as a conscious process and this can be a source of significant errors. Analysis of gingival composition is also a determining issue in positioning the collar of the lateral incisors in a location that is aesthetically optimal. The gull-wing profile, where the collar of the lateral incisors is slightly more coronal than that of the front teeth or the canine teeth, is considered to be more attractive according to Chiche⁵ (Fig. 2).

Dental aesthetics

As regards dental aesthetics, the proportions of the proposed implant supported tooth can reflect two different scenarios:

Fig. 4: According to Levin, following the golden ratio, the width of the lateral incisor $y = 0.62 x$ and, for Preston, it is $0.66 x$ (images from Papathanassiou).⁶

Fig. 5: Evidence of bone deficit at 22 (case shown in Fig.1).⁶




Fig. 6




Fig. 7




Fig. 8


Fig. 6: Simulation of location of 3mm⁶ implant in cross section (case shown in Fig.1).⁶

Fig. 7: Evidence of radicular convergence.

Fig. 8: Orthodontic layout of implant corridor.

Fig. 9: Diastemas created around a riziform tooth to obtain a space of 6 mm.

Figs. 10a & b: Centered location of zenith of 22 (**a**) (arrow) to be taken into account when making the crown 12 (**b**).



- there is a unilateral missing tooth and the contralateral incisor has normal and aesthetically pleasing proportions. The objective will be to create a lateral incisor implant that is a mirror image;
- with the same situation but where the contralateral incisor is small; this is a situation that occurs frequently in unilateral agenesis where the incisor that is present is riziform or, if there is agenesis of both lateral incisors, the clinical examination should gather the information required to decide on the dimensions and coronal axes of the proposed lateral incisors. An analysis of the occlusion and the dimensions of the central incisors are the clinical parameters that make it possible to establish the characteristics of the planned prosthetic teeth.

The anatomy of the lateral incisor has been the subject of various publications, including, notably, by Papathanassiou⁶ who defined average dimensions and a typical form (Fig. 3a) and also presented numerous morphological variants affecting these dimensions and also other characteristics such as the crown/root ratio and the coronal and root axes (Fig. 3b). These morphological criteria, which can now be found using 3-D imaging, have had a significant influence on the location of implants in all spatial planes in order to achieve the goal of harmony of form and dimension. Other publications, such as those by Levin⁷ and Preston,⁸ make it pos-

sible to estimate the width of absent lateral incisors on the basis of the central incisors (Fig. 4).

“Establish the **ideal width and orientation** of the planned prosthetic crown.”

Implant location

A clinically significant deficit signals the need for reconstruction of hard tissue but, conversely, a site without a tooth with no loss of volume should be subjected to a three-dimensional X-ray, as thick soft tissue can hide a lack of hard tissue (Fig. 5). A thin tissue biotype or a lack of attached gingiva can be a sign that gingival augmentation surgery will be required, particularly if a bone graft needs to be performed.

Occlusion

For orthodontic treatments, the anterior guidance should be analyzed carefully. It can be tempting to increase the perimeter of the maxillary arcade in order to obtain, at the least, implant corridors that are sufficiently wide at the level of 12 or 22. However, an overjet will make it very likely that the natural teeth will move in relation to the implant prosthesis with highly negative consequences for the sustainability of the cosmetic outcome.

Documentation

Taking photographs at the start of the treatment will make it possible to maintain a record of the initial condition, which is always useful if there are medical/legal problems at the end of the treatment. In addition, the images often make it possible to see problems relating to width, axis or asymmetry that sometimes go unnoticed during a clinical examination.

“Check **anterior guidance and absence** of overjet.”

Complementary tests

2-D imaging

Panoramic X-rays or retroalveolar radiography make it possible to check the depth of implantable




Fig. 11: Cortical graft in place, shaped to support future papillae (case as shown in Fig. 1).

Fig. 12: X-ray result, compare with Fig. 6.

Fig. 13: Clinical outcome five months after the graft. Compare with Fig. 11.

Fig. 14: 3 mm diameter NobelActive implant.

bone in relation to the floor of the nasal cavity, the bone level in relation to that of adjacent teeth and the parallelism of the central incisor and canine.

3-D imaging

3-D imaging is required to check the vestibular palatal dimensions of the bone crest. There are three possibilities:

- the crest is sufficiently wide to take an implant without any bone augmentation;
 - the crest is narrow, bone augmentation is required prior to siting the implant (Fig. 6);
 - intermediate situations where the siting of the implant will be accompanied either by bone splitting or by guided bone regeneration.
- Orthodontic preparation

When the adjacent teeth present apical convergence, the orthodontic preparation should create a mesio-distal dimension at the level of the root that allows the implant to pass with a margin of at least 1 mm of bone (Figs. 7 & 8). Where there is a contralateral incisor of a normal size, the rule for the orthodontist is to measure the width of that tooth carefully and to recreate the same width in the crown of the planned implant. Where the contralateral incisor is riziform, the orthodontist should plan the future face of the tooth in order to achieve two laterals with the same shape.

Diastemas around the riziform tooth make it possible to achieve a smile that, in the end, is almost symmetrical (Fig. 9). The riziform incisor does not have to be in the centre of the space but should be positioned in such a way that the papillae and the future zenith of the tooth are optimized. The zenith should be located 0.4 mm distal from the centre of the tooth for a lateral incisor, according to Chu et al.⁹ (Figs. 10a & b). Sometimes, a zenith situated more than 1 mm from a line between the collars of the central incisor and the canine should be surgically altered by coronal lengthening as a lateral incisor that is too short can also be aesthetically unacceptable.

“The orthodontist should **anticipate the future prosthetic morphology of the riziform incisor.**”

Hard tissue augmentation

Where a bone reconstruction is indicated, this should take into account one of the key factors for the overall cosmetic outcome: restoration of papillary support in order to avoid any unsightly black triangles between the lateral incisor and the adjacent teeth or any concave area above the implant crown that would create an ugly shadow.

The cortical graft, taken from the chin or the external oblique, should be formed in such a way as to provide support for the gingival papillae (Fig. 11). Gaps under and around the graft should be filled with cortical bone particles, crushed from the chin block or lateral mandibular area using a bone mill.

The attachment must be reliable. This is done using two 1.6 mm diameter osteosynthesis screws (Fig. 12). Autografts take about 5 months to heal.

Figs. 15a & b: Papillary view (b), X-ray view (a), two years after the insertion of the implant.






Fig. 16




Fig. 17




Fig. 18



Fig. 19




Fig. 20




Fig. 21

Fig. 16: Mobile Anyray 2 (VATECH) X-ray generator.

Fig. 17: Intraoperative X-ray, Precision Drill inlay (left on picture) and in situ (right on picture).

Fig. 18: Clinical outcome five months after the graft. Postoperative X-ray NobelActive 3/0 implant and 15° abutment in place.

Fig. 19: Initial situation.

Fig. 20: De-epithelialisation of a palatal flap into a diamond shape.

Fig. 21: Unfoldment of palatal flap, vestibular edge.

Ideally, the implant should be inserted between 4.5 and 5.5 months after the graft (Fig. 13).

“Fully reconstruct papillary support.”

Insertion of implant

Choice of implant

The mesio-distal dimension of the gap will determine the choice of the implant. When this is close to or less than the average size of 6.5 mm, the bone and papillary volume around standard size implants will be limited. According to Hasan et al.¹⁰ and Bourauel et al.,¹¹ the disadvantage of small diameter

implants is that they transmit higher stresses to the crestal bone than do standard implants. When replacing a lateral maxillary incisor, it is possible to arrange both the anterior guidance and the deduction in such a way as to make them largely affect the natural teeth, in the absence of any significant malpositioning, and in this way reduce the stresses applied to the implants. Under these conditions, small diameter implants have the advantage of increasing surrounding residual bone volume as well as space available for papillary healing.

In a forthcoming study of 120 NobelActive 3 mm diameter implants, one of the conclusions confirmed the importance of these small diameter implants as regards the additional height of the papillae, resulting in an improvement in the Fürhauser pink aesthetic score¹² (Figs. 14, 15a & b).

Fig. 22: Creation of flap envelope, Swann-Morton blade through the envelope. SM 63, inlaid with transparency.

Fig. 23: Passage of suture through the envelope.

Fig. 24: The palatal flap is folded into the vestibular envelope using a suture thread.



Fig. 22




Fig. 23




Fig. 24



Fig. 25



Fig. 26

Fig. 25: Tissue integration with ceramic crown.

Fig. 26: Vestibular bulge obtained with modified flap.

“Favor **small diameter** implants.”

3-D positioning

As regards replacement of a lateral maxillary incisor, the tolerances for the location of the implant are very small because of the narrow width of the implant corridor. Two recent meta-analysis^{13,14} concerning the precision of surgical guides resulting from 3-D imagery, even if these do not apply specifically to the lateral incisor replacement, has found a deviation in the order of a millimetre at

the point the implant emerges and 4 to 5 degrees as regards the drilling axis. For Van Assche et al.,¹⁴ the average imprecision at the apex of the implant is 1.24 mm.

Since these measurements are incompatible with a 12 or 22 implant corridor, it is important to check the first drill hole(s) during the operation, whether the surgery is guided or being carried out freehand. If the implant clinic does not have retro-alveolar X-ray equipment, portable generators such as the AnyRay II (VATECH) are available on the market, which allow you to produce intraoperative images (Fig. 16).

AD



PUBLISH
YOUR
EXPERTISE!

Become an author for implants—
international magazine of oral
implantology

Please contact:

Georg Isbaner
Editorial manager
g.isbaner@oemus-media.de





Figs. 27a–c: Clinical and X-ray views, vitroc ceramic in place.

In this context the Precision Drill from the Nobel Biocare kits is particularly helpful. Its sharp point provides considerable precision at the point of entry and its small dimensions make it possible to correct any deviations from the ideal axis occurring during the first drilling (Fig. 17).

In the vestibular palatal plane, it is essential to prepare a prosthetic treatment plan before inserting the implant because the positioning requirements differ:

- for a screwed prosthesis, the axis of the implant is very strictly determined by the point in the cingulum where the screw emerges;
- with a cemented prosthesis, the tolerance is slightly greater as it is possible to make a correction to the axis by an abutment angled up to 15 degrees or by a Procera type individualized abutment (Fig. 18).

“Position the implant under **X-ray** monitoring.”

Soft tissue management

Whether the soft tissue management is carried out at the time the implant is put in place or when it is exposed, the choice of surgical technique depends on an examination of the initial situation:

- horizontal deficit of soft tissue that could result in the underlying titanium being visible;
- vertical deficit in the papillae that could result in unsightly black triangles.



Different surgical techniques can be used, depending on these deficits, which are taken from three publications: the roll flap developed by Abrams,¹⁵ the envelope technique of Peter Raetzke¹⁶ and Carl Misch's split-finger.¹⁷

- if there is just a horizontal deficit, a modified rolled flap⁶ can be carried out, without separation of papillae and without vestibular incisions, the palatal flap being folded into an envelope flap (Figs. 19 to 25). The attraction of this technique for the patient is that a second operation site to take a graft is not required. In addition, it makes it possible to recreate a root eminence, considered already 20 years ago by Silverstein and Lefkove¹⁸ to be an important factor for the aesthetic outcome (Figs. 26 & 27a to c);
- where there is a vertical deficit, a crestal W-shaped incision as described by Carl Misch¹⁷ is indicated. This makes it possible to recreate

Fig. 28: Initial incision creating two vestibular half papillae.

Fig. 29: Suture of half papillae (situation in Fig. 1).

Fig. 30: De-epithelialisation tuberosity graft.






Fig. 31



Fig. 32

Fig. 31: Insertion of connective graft buried under the papillae.

Fig. 32: Clinical outcome in a case of gummy smile.

an anatomical gingival architecture while, as a first step, creating two vestibular neo-papillae (Fig. 28). After separating the sections, the palatal tissue (finger) is divided into two to make two palatal half-papillae, joined one on one with their vestibular counterparts (Fig. 29);

- where there is a combined deficit, the same incisions are combined with a buried connective vestibular graft. Provided that there is sufficient volume, the graft is taken from the maxillary tuberosity, since this area has the advantage of providing graft tissue that is more dense, opaque and less adipose than the palate and, in addition, results in less postoperative pain. If the graft is transferred in a V- or Y-shape, it can support the newly formed papillae. The shape of the palatal incision can be modified to a Y-shape to assist rotation of the palatal half-papillae (Fig. 31).

If the thickness of the buccal gingival tissues has not been augmented or if collagen substitutes are used that do not have the opacity characteristics of tuberosity connective tissue, the aesthetic outcome can be compromised. If there is recession of the external table or the titanium abutment under thin connective tissue, the grey titanium colour can be seen through the gum as a grey halo above the crown collar, which is detrimental to the aesthetic appearance (Figs. 32 & 33).

of hard or soft tissues, will result in aesthetic problems. The prosthetic stages allow optimisation of the result as regards the gingival context but any error in the surgical stage will often be impossible to correct during the prosthetic stages. For this reason it is vital to approach this first part of the implant treatment for a lateral incisor with thoroughness and precision.



Fig. 33

Fig. 33: Insufficient soft tissue thickness alters the chromatic outcome.

Editorial note: A list of references is available from the publisher.

This article was originally published in the Éditions CdP prosthetic journal, No 167, September 2014 and the Clinical Masters Magazine No 1/2015.

“Systematically augment the thickness of buccal connective tissue.”

Conclusion

The aesthetic fundamentals for an implant are in the preprosthetic surgical stages of the treatment. Any approximation in the location of the implant in such a narrow implant corridor, any lack of support for papillae or any deficiency in the thickness

contact



Dr Philippe Russe,
former assistant at the
Reims University and Hospital.
He runs his private practice in
Reims, France.

Dr Patrick Limbour, MCU-PH, Head of department of oral surgery, Pontchaillou University Hospital, Rennes, France.