


Integration of diode laser surface decontamination in periimplantitis therapy—a twelve year review of a fit for practice concept

author_ Georg Bach, Germany

Manifestation of periimplantitis
 On probing, secretion is released at the mesial implant, though the clinical appearance is inconspicuous and further probing leads to a substantial bleeding. After mobilization of the soft tissues, the typical crater-shaped periimplant bone defect becomes visible.

After many years of great euphoria, a certain disillusion has spread in implantology, which is especially due to the reason that implants with corresponding suprastructures do not last forever, like it has often been pointed out. Anyway, complications cannot totally be excluded. Professor Herbert Deppe, Chair for the Dental Surgery and Implantology Department of Munich University, has recently reported on the fact that approximately an eighth of incorporated implants show periimplantary lesions after about 10 years. In the beginning, the main fear was that enossal implants had to face early complications. Nowadays, this is no more the case since sophisticated surgery techniques and improved implant surfaces have reduced these risks. One still has to worry about long-term sequelae shown in artificial abutments caused by periimplantary lesions af-

ter some years of strain. However, periimplantitis is mainly induced by bad oral hygiene and/or the inability to carry out mouth care (e.g. in old patients), and it is not associated to a certain type of implant (system-independent). Numerous therapy approaches have been made to preserve artificial abutments suffering from periimplantitis. A four phase treatment model is usually applied (hygienization phase, surgical resective phase, reconstructive and augmentative phase, recall phase). This model has considerably been enhanced by the launch of diode or injection lasers, which have later been complemented by CO₂- and Er:YAG-, and Er,Cr:YSGG-laser respectively. Since the mid-nineties, diode lasers belong to the established wavelengths used in dentistry. Today, diode lasers with short pulse technique are predominant,



In Kooperation mit

Gothaer / AMG

Stiftung | Mensch & Medizin®

Neu!

- 3,5 mm Implantatdurchmesser
- Garantieverlängerung 200,00 Euro* Festkostenzuschuss zur Prothetik
- deutschlandweite Medienpräsenz für Lizenzpartner

Jetzt!

Periointegration® ...statt Periimplantitis

In Kooperation mit dem Fraunhofer Institut und namhaften deutschen Universitäten wurde eine intensive Grundlagenforschung im Bereich moderner Hartstoffschichten betrieben. Ein Technologietransfer aus der Luft- und Raumfahrttechnik ermöglicht erstmalig die Beschichtung von Titanoberflächen mit Zirkoniumtrioxid und Zirkoniumoxid.

Die aktuellen wissenschaftlichen Ergebnisse auf dem Gebiet der Oberflächentechnologie zeigen, dass Zirkoniumtrioxid ein Anhaften des Biofilms mit parodontalpathogenen Keimen erheblich verringert und die Anlagerung der Gingiva zu einem dichtanliegenden Saumepithel maßgeblich fördert. (Größner-Schreiber et al. 2006).

Periointegration bedeutet die langfristige und entzündungsfreie Erhaltung von unterstützendem Knochen mit einer dicht anliegenden Weichgewebssmanschette um das Implantat.

In Zusammenarbeit mit der Academy of Periointegration, ein Zusammenschluss führender Vertreter aus Forschung, Lehre und Praxis, ist es Clinical House Europe GmbH gelungen, sieben Designkriterien in eine neuartige ultradichte, periointegrative Implantatkonstruktion umzusetzen.



Vollkaskoimplantat® ...wir tun was

Das Vollkaskoimplantat ist das erste Business Erfolgskonzept für die Implantologie, welches Technologie, Garantieabsicherung und Medialeistung zu einem bundesweit erkennbaren Markenkonzept verbindet. Informieren sie sich über die Vorteile für Ihre Praxis und die Zugangsvoraussetzungen für Lizenzpartner.

Tel: +49 (0) 234 90 10 260

info@vollkaskoimplantat.de

www.vollkaskoimplantat.de

Clinical House Dental GmbH

Am Bergbaumuseum · 44791 Bochum

Tel. +49 (0)234 90 10 260 · Fax +49 (0)234 90 10 262

Gratis!

Vollkaskoservice

7 Gründe für das Lizenzkonzept Vollkaskoimplantat:

- ✓ wissenschaftlich dokumentierte Periointegration®
- ✓ erstes Business-Erfolgsrezept für die Implantologie
- ✓ 10 Jahre Recall-Motivation für Patienten
- ✓ Gothaer/AMG abgesicherte 10 Jahres Garantie*
- ✓ Festkostenzuschuss Prothetik von 200 Euro*
- ✓ Chirurgische Kostenerstattung*
- ✓ deutschlandweite Medien-Präsenz

Ja bitte senden Sie mir Informationen zum Thema **Vollkasko Infogutschein**
 Abchirma bitte faxen an (0234 90) 0262. Oder per Post an Clinical House Dental,
 Am Bergbaumuseum 31, 44791 Bochum.

Name _____

Anschrift: _____

Tel: _____

Email: _____

www.stiftung-mensch-und-medizin.de
www.vollkaskoimplantat.de

*gemäß AGB's

Foto: corbis

though it all started out with the cw mode. High performance diode lasers emit monochromatic, coherent light of wavelength 810 nm, which is especially well absorbed by dark surfaces. Thanks to these physical conditions, the injection laser (= diode laser) is perfectly suitable for incisions applied in standard dental surgery, as well as for the resection of benign tumors in the oral cavity, the uncovering of implants and for application in mucogingival surgery. The good cutting properties of diode lasers are due to the extraordinary absorption of laser light by the hemoglobin located inside the tissue. Additional to soft tissue surgery, the diode laser is also used for decontamination of surfaces covered with microbes (on implants and teeth). It could be demonstrated that especially the Gram-negative, anaerobe microbiological spectrum was properly damaged by laser light (Bach und Krekeler (1995;2000)). In compliance with reasonable performance and time parameters, which have been confirmed sustainably by clinical long term studies (Moritz (1996), Gutknecht (1997), Bach et. al. (1995, 1996, 1998, 2000, 2001)), a thermic or morphological damage of the implant surface and the surrounding bone tissue can definitively be excluded (Bach and Schmelzeisen (2002)). It was the aim of the present study to demonstrate and evaluate a treatment model for periimplantitis therapy, which shows sustainable results and which is absolutely suitable for practice. There is no doubt that the conventional methods for periimplantitis treatment, which have often been described in literature, permit adequate surface cleaning and thus also the reduction of pathogenic microorganisms on the implant surfaces. Nevertheless, the complete removal of relevant bacteria cannot be ensured. Moreover, the conventional removal of biofilms has only little influence on those bacteria infiltrating the soft tissue. The integration of diode laser light in periimplantitis therapy must be seen as a new approach.

_Material and method

Ten patients (with n = 17 implants) have been treated and examined for a period of more than 12 years (since May 2007). In spring 1995, all of them suffered from periimplantitis on their artificial titanium abutments.

_Pathogenesis of periimplantitis

Periimplantitis therapy represents a border area between implantology and parodontology. The causes for parodontitis and periimplantitis are bacterial infections, in particular they are biofilm based infectious diseases. Gram-negative and anaerobe microbes are mainly responsible for the destruction of the parodontal and periimplantary supporting tissue. As a rule, one of the following microbes

causes parodontopathy in case of one of both biofilm based infectious diseases:

- _ *Actinobacillus actinomycetemcomitans*
- _ *Prevotella intermedia* and
- _ *Porphyromonas gingivalis*

Whereas periimplantitis is mainly caused by the following microbes:

- _ *Fusobacteria*
- _ *Prevotella intermedia* and
- _ *Porphyromonas gingivalis*

The principal object of periimplantitis therapy carried out in our dental clinic was to remove the biofilm and hence the removal of the mentioned pathogenic microorganisms.

_ Patients treated

For detailed data, age and sex of the patients, please see Figs. 1 and 2. It should be mentioned that an accumulation of the disease's first incidence is registered in the middle years (age: 30 to 50 years) in both groups. Sex-specific differences could not be ascertained.

| Age | Number of patients |
|-------------|--------------------|
| 20–30 years | 1 |
| 30–40 years | 3 |
| 40–50 years | 3 |
| 50–60 years | 2 |
| 60–70 years | |

Fig. 1_ Age pattern of the examined and treated patients in 1995.

| Sex | Number of patients |
|--------|--------------------|
| Female | 5 |
| Male | 5 |

Fig. 2_ Evaluation according to the sex of the examined and treated patients.

_Inclusion and exclusion criteria

All patients involved had to meet strict inclusion criteria as there were:

- _ Clinically visible inflammatory signs like BOP (bleeding on probing) and high probing depths
- _ Radiovisible periimplantary bone lesions ("crater")

Exclusion criteria were:

- a) Severe primary diseases
- b) Nicotine or alcohol abuse
- c) Lack of compliance

Due to the strict inclusion and exclusion criteria only a limited number of people could be admitted for this study.

_Treatment procedure

Equal treatment procedures for all periimplantitis patients:

39TH INTERNATIONAL ANNUAL CONGRESS OF THE DGZI

Complications, Risks and Failures in the Oral Implantology

October 9–10, 2009, in Munich
Hilton Munich Park


SPECIAL PODIUM

„Periimplantitis –
a challenge of
the future“ with
talk of experts



Allianz Arena

Information:
Phone: +49-3 41/4 84 74-3 08
Fax: +49-3 41/4 84 74-2 90
event@oemus-media.de
www.dgzi.de
www.event-dgzi.de



Periimplantitis therapy:

You can see the first patient, who had undergone periimplantitis treatment, by means of diode laser decontamination, according to our model.

November 1994: Manifestation of periimplantitis at implant regio 13. The panoramic tomography (detail) shows a significant bone loss at the artificial abutment. After mobilization of the soft tissue the situation of the defect becomes clearly visible.

January 2008: The prosthesis made in 1990, is still in the same position. The situation of the treated regio 13 implant does not show any irritations with and without suprastructure.

There is no evidence of probing depth. The panoramic tomography shows a stable bone situation. Besides the reconstructed defect of regio 13, only the root filling of 43 protrudes. This is the only difference compared to the tomography taken in 1995.

1. Initial therapy:

- _ Motivation and instruction of patients
- _ Cleaning and polishing
- _ Application of disinfecting agents

2. Resective phase:

- _ Forming of a mucoperiosteal flap
- _ Removal of granulation tissue
- _ Decontamination by means of diode laser light ($p = 1.0$ watt, $t_{max} = 20$ sec.)
- _ Apical shifting of soft tissues

3. Reconstructive phase:

- _ If necessary, bone augmentation
- _ Where applicable, mucogingival corrections

4. Recall phase:

- _ After four weeks, six months, one year and then annual evaluations of clinical findings, taking of X-rays (PSA), decontamination of eventually exposed areas by means of diode laser light.


_Image processing methods

As a rule orthopantomograms (panoramic tomography) and additionally dental films in parallel technique were chosen as an adequate image processing method. In some cases of exacerbated in-

flammations A/B scan ultrasonic methods were applied. A preoperative orthopantomogram and the dental film status (dental shots of the respective areas) were taken. A postoperative orthopantomogram was directly taken after surgery. A panoramic tomography was taken one year later and then every two years. The advantage of the orthopantomogram is its panoramic-like view of all teeth, the osseous limbus alveolaris and important neighboring anatomical structures. The dental film in parallel technique allows statements concerning progredience, stagnation of loss of hard- and soft tissue, and it shows the course of the limbus alveolaris in a reproducible way.

_Microbiological diagnosis

Time schedule: Preoperative, four weeks postoperative, one year postoperative and in a five to ten-year postoperative interval germs were eliminated from the effected areas. We did not apply the classical microbiological examination technique (isolation of microbes—cultivation – pure cultures – microscopic samples – gas chromatography – antibiotic sensitivity testing—and biochemical identification, the so called "bunte Reihen"). We used DNA-RNA hybridization probes instead. The advantage of these hybridization probes is that no living



Saving of a prosthesis by treating periimplantitis of a strategically important implant in the upper jaw.

March 1995: Just one year after the incorporation of a very sophisticated and for the patient nearly too expensive implant-supported prosthesis in the upper jaw, the manifestation of periimplantitis was detected in the first quadrant. After mobilization of the soft tissue (below) the defect situation becomes clearly visible. Four months after the surgical resective phase there were no clinical signs of irritation.

November 2007: The prosthesis is still in its intraoral place. Meanwhile, the patient has reached the age of 63 years. The situation of the treated implant regio 13 (total suprastructure) does not show any clinical signs of irritation in toto and in the former surgical area. There is no probing depth.

material of the areas probed is needed for cultivation purposes, which minimized the work in the dental clinic (without direct access to an Institute of Microbiology). Additionally, the results were much faster on hand as is the case with classical microbiological examinations. The disadvantage of this rapid test is its high price. Furthermore, only special marker microbes can be detected and not all pocket microorganisms can be determined. The germ extraction site had to be dried carefully with a cotton swab, the paper tip was placed, and after a waiting time of 10 seconds put into a sterile storage vessel and sent to the manufacturing company for microbiological diagnosis. The company is in charge of microbiological diagnosis and evaluation of the so called microbe marker values. The classification of marked microbes was: less than 0.1 % = negative; 0.1-0.99 % = low; 1.0-9.9 % = middle, more than 10 % = high.

_Laser light decontamination

Decontamination formed an essential part of the whole therapy. It was carried out by means of diode laser light with 1 watt performance and 20 seconds of application time per implant under fiber contact. A special program (I = implantology-parodontology) was at our disposal, which was used together with the corresponding device (Oralia 01 IST). Performance and time limitation (1.0 watt, 20 seconds) were already fixed parameters of this program. When observing these parameters (time limitation and limitation of performance) it can be guaranteed that the disease causing microbes will be damaged sufficiently and thus, pulpa, periimplantary and periodontal tissue structures will not suffer any thermic damages (Bach and Krekeler (1995)).

_Results

Alltogether 10 patients could be examined and checked up during the whole 12 years. In 1994/1995 the "Diode Laser Basic Study" of the Department of Periodontal Surgery of the Dental Clinic in Freiburg/Germany included 50 periimplantitis patients. Due to moving, change of dentist, dead of pa-

tients and other unknown reasons the number of patients was reduced to 10, who are still patients of my dental clinic.

a) Microbiological results

For microbiological results please see Fig. 3. It must especially be emphasized that Porphyromonas gingivalis could nearly be completely eliminated during the whole examination period, and a significant reduction of other anaerobe, Gram-negative bacteria could be achieved. We could obtain similar results for Porphyromonas gingivalis and Fusobacteria except for two cases of low concentration and one of middle concentration, these bacteria could be limited to the lower level of detection in other patients, whereas other relevant marker microbes could be considerably repressed.

b) Recurrence

One of the following results was considered to be a case of recurrence:

- _ Occurrence of probing depths of more than 4 mm
- _ Loss of implant
- _ Recurrence of an inflammation
- _ Excessive soft tissue inflammation with pocket activity

After 12 years the quota of recurrence was 23% in the periimplantitis group (4 implants). It is stated in international literature that the five year observation period recurrence rate is 30 %.

c) Losses after 140 months

Within the examination period of 12 years we suffered the following losses: two of 17 implants (12 %).

d) Radiological results

On the occasion of the one year check up, a reconstruction of the once crater-shaped defect could be found at the first thread and implant cervix respectively in all 17 implants. After five years this was the case in twelve implants, after ten years in ten implants and in nine implants, when the last X-ray control was carried out. In two implants a successive loss of the bony supporting tissue forced us to remove the artificial abutment in one case af-