

# Single-tooth replacement using CBCT matching and virtual wax-up

Dr Jakob Zwaan, Netherlands & DT Vito Minutolo, Italy

**In our practice we often** encounter major challenges when just a single tooth needs to be substituted. In order to estimate the risk of an unacceptable aesthetic treatment result and to determine the most effective treatment plan, it is necessary to perform an analysis of the desired tooth shape, the soft tissue architecture and the bone volume necessary to stabilise an implant in the optimal position and support the soft tissue.

In the following, an approach to cases will be described in which a single tooth needs to be replaced by an implant-supported crown. After an anamnestic interview with the patient, we proceed with the intra-oral examination. Hygiene and periodontal health are checked. Normally, an intra-oral radiograph for a single tooth is taken first (Fig. 1) or a dental panoramic tomogram if the need for a more extensive treatment is suspected. In the same session, both dental arches are scanned with an intra-oral scanner and the bite is registered. The photographic sequence is as follows:


1. Full frontal view intra-oral photograph.
2. Detailed photograph of the single arch, possibly with a black mirror to contrast the teeth (Fig. 2).
3. Photograph of laterolateral detail of the tooth and gingival profile (Fig. 3).
4. Full-face photograph with maximum gingival exposure.
5. Full-face photograph of a spontaneous smile.
6. Photograph of the full face at rest.







This sequence allows one to view immediately the presence of orthognathic and periodontal issues (Fig. 1), to evaluate the biotype (Fig. 2) and to estimate aesthetic challenges, like tooth colour, tooth texture, soft tissue/lip exposure and position of the incisal edge/lip (Fig. 3). The 3D intra-oral scan is helpful for determining orthodontic alignment of the teeth and in our protocol replaces an occlusal and/or 12 o'clock photograph in most cases. "There can be different ways of treating a disease, but there can be only one correct diagnosis." Dr Morton Amsterdam, 1974. When anamnesis, intra-oral examination and preliminary radiographs conclude that the tooth in question cannot be preserved, it needs to be decided what the optimal timing for extraction and a CBCT scan is and how to provide for a temporary tooth replacement. Also, the operator must choose between immediate, early or delayed placement in the fresh extraction socket.

Our policy is the following: in case of acute inflammation that cannot be effectively treated, we will proceed with extraction. A temporary fixed etch and bond or removable prosthesis can be used to guarantee acceptable aesthetic comfort to the patient. In these cases, a CBCT scan will be taken after extraction so that the most detailed image of the socket anatomy can be obtained. Since a provisional solution has been provided for, there is no need for very early implant placement. Timing is now based on the expected period needed for the infection to be eliminated and the risk of loss of volume by the collapse of tissue. Normally, the implant is placed four to six weeks after the extraction. Another reason for delayed implant placement can be the need for healed soft tissue in order to facilitate proper wound closure to protect, for example, bone substitutes and membranes when bone augmentation is necessary.

If the patient is suffering, extraction should be done quickly. If the anatomy and biological conditions are favourable, implant surgery can be carried out as early as one week after extraction. Only if there is no inflammation or infection, and sufficient bone and soft tissue quantity and quality are present, the implant should be placed in the fresh extraction socket. In such a case, the CBCT scan would be performed before proceeding.





Minor bone augmentation and/or connective tissue grafting can be performed contemporaneously. The decision to place an immediate provisional crown on the implant is strongly related to the expected primary stability of the implant, as well as the opportunity to manage the position of biomaterials in such way that undisturbed and uncontaminated healing is guaranteed. After healing, good aesthetics and sufficient protection of the underlying implant and implant–prosthesis connection are requisite if we wish to treat our patients in the best possible way.

## Risk evaluation

### First aesthetic risk evaluation

One can start with a render of a 2D photograph. We use the macro intra-oral shot with the black background behind the teeth (Fig. 2). With Adobe Photoshop, GIMP, Microsoft PowerPoint or Keynote, for example, it is easily possible to cut out the shape of the contralateral tooth that will not be extracted, copy it, flip it horizontally and paste it in the position of the tooth that needs replacement. It will be clear immediately whether this shape supports the papillae sufficiently or whether there is a lack of volume that needs to be compensated for. Also, this image with the flipped contralateral tooth can be aligned with the original photograph and then draw a horizontal line across both images that coincides with the same gingival reference points. This will demonstrate whether there is a vertical component that indicates a lack or abundance of soft tissue. This can be easily quantified in a metric system if an intra-oral reference is measured with

a calliper. The patient can now be informed whether additional procedures like guided bone regeneration (GBR) or a connective tissue graft will be needed.


### Second risk evaluation

The intra-oral scan is imported into CAD software and transformed into a virtual master model without the tooth to be extracted and a separate STL shape of the ideal CAD-designed tooth. Now there is the opportunity for 3D evaluation of the dimensional relation between the new tooth and the soft tissue before extraction. In the current case, the tooth involved had not been extracted and a CBCT scan was performed (X-Mind trium, ACTEON; 110 x 80 mm field of view; 0.15 mm voxel size) for further investigation and treatment planning. In the AIS 3D App software that comes with the CBCT X-Mind trium device, STL files can be matched and aligned with the 3D bone volume, thus giving the opportunity to plan the future implant position taking into account the shape and position of the future crown (Figs. 4a & b). In accordance with the prosthetic procedure preferred, cemented versus screw-retained, CAD/CAM-fabricated versus manual layering and the type of material to be used, all the information for the final treatment plan is available, on which decisions can be made regarding GBR, connective tissue graft and timing of implant loading.

## Case report

The female patient, aged 47 and a non-smoker, was in good general health. She experienced increasing mo-





bility of the maxillary left central incisor and complained about compromised aesthetics due to the extrusion and progressive migration of the tooth in a buccal direction. The incisor had been treated with a crown at a preadolescent age after a violent trauma. The intra-oral radiograph showed incomplete root development and evidence of a root canal therapy suggesting a strip perforation though no signs of periapical lesions were present. The shape of the crown was not symmetrical in relation to the triangular shape of the maxillary right central incisor, but had a wider and rectangular profile. Minor general gingival recession had led to the presence of a tiny interdental space. The marginal gingiva was reddened, and the central papilla was not symmetrical. Probing depths were within 2mm for both the right and left central incisors and the radiographic mesial and distal bone peaks were of a regular height. The photographic aesthetic evaluation showed that it would be very difficult to obtain symmetry in tooth shape and have good-looking and healthy soft tissue support at the same time. The patient's maximum smile exposed the gingival contours. In such cases, it may be wise to consider also the possibility of altering the anatomy of the contralateral tooth with, for example, a ceramic veneer and discuss outcomes with the patient before finalising the treatment plan. This can be evalu-

ated by performing the cut/copy/flip/paste sequence in reverse. It was decided to start performing the best possible replacement of the maxillary left central incisor and evaluate at an advanced stage with a temporary crown on the implant and mature, conditioned tissue whether to add a veneer to the maxillary right central incisor.

### Analysing the CBCT scan

The short-rooted tooth could be extracted without compromising the buccal bone, and there was sufficient bone volume and quality to obtain good primary implant stability. Thanks to the AIS 3D App software, this information can be visualised using the bone density tool and linear measures tool (Fig. 4c). The presence of the nasopalatine duct prohibited ideal palatal positioning of the implant, and if the implant were to be placed flush with the palatal alveolar bone, this would have resulted in a 1.5–2.0mm high exposure of the implant collar on the buccal aspect (Fig. 5a). It was thus decided to place the implant in that position and to augment the buccal bone volume with a contemporaneous GBR procedure, thus also providing for major soft tissue support. Literature suggests that that in some measure the implant will deviate buccally from the orig-






Fig. 16




Fig. 17

inal planning because of the major mechanical resistance of the palatal plate.<sup>2-4</sup>

The author's team prefers screw-retained solutions. Several production centres are capable of milling angulated screw access holes in cobalt-chromium abutments of up to 25°. It can be easily checked in the implant planning software whether the future access hole will exit on the palatal aspect of the tooth, either by angulating the implant extension tool or by choosing a virtual abutment from the library. Confirming being in the safety range from this point of view allowed for an approach that foresaw the implant in native bone without the necessity for major GBR on the apical aspect of the implant. It was decided to use a surgical guide (Figs. 5a & b) for only the first drill to determine with precision the position and angulation of the osteotomy that would be performed freehand thereafter. A removable temporary prosthetic tooth was produced in advance.

### Surgery

Local anaesthesia was performed with 2% mepivacaine with 1:100,000 adrenaline. Preventative antibiotic therapy with amoxicillin (1 g, b.d. for five days) was prescribed,

aided by use of a 0.2% chlorhexidine mouth-rinse three times a day. The tooth was extracted and the sulcular epithelium removed with diamond burs. The milled surgical template (Figs. 6 & 7) served as a guide for the first 2 mm diameter pilot drill (Fig. 8). Thus, the planned depth, position and angulation of the osteotomy were obtained. The drill sequence was completed freehand, using tapered 3.0 and 3.4 mm drills. A Neoss Pro-Active Tapered Implant of 4 mm in diameter and 13 mm in length was inserted flush with the mesial/palatal/distal bone, motor driven up to a torque of 50 Ncm and then with a manual wrench (Fig. 9). The correct position of the internal hex was verified by checking the references on the implant driver, which ideally points in the buccal direction. Resonance frequency analysis with Penguin-RFA (Integration Diagnostics Sweden) determined an ISQ value of 73/76. A Neoss Esthetic Healing Abutment with a ScanPeg was connected to the implant (Fig. 10). A flap was then raised after a vertical incision of the frenulum and the expected buccal exposure of the implant neck was evident. Autogenous bone was positioned directly on the implant surface (Fig. 11), followed by a bone substitute on top of it and on the buccal cortical bone (Fig. 12). This material was covered with a resorbable membrane (Fig. 13). The mobilised flap was then repositioned by rotating it coronally and fixed with single su-




Fig. 18



Fig. 19






Fig. 20




Fig. 21




Fig. 22

Fig. 23




Fig. 24




Fig. 25

tures (Fig. 14). The removable partial denture was adapted and delivered (Fig. 15). An immediate postoperative CBCT scan of 60 x 60mm was performed, and it confirmed a perfectly centred implant position (Figs. 16 & 17).

### Intra-oral scan

Eight days after surgery, the patient reported that healing was uneventful and the prosthodontist removed the stitches. It has become the author's standard protocol to perform an intra-oral scan for implant position in this same session (Fig. 18). The specific and unique PEEK healing abutment used has an internal circular channel and on one side, normally positioned on the buccal aspect, a vertical rectangular slot (Fig. 19). After removing the PTFE tape used to plug this area during surgery, a ScanPeg can be positioned inside the healing abutment. This allows for a unique scanning procedure without removing the healing abutment, thus avoiding disturbing healing tissue or dislocating recently placed biomaterials.

### Temporary crown

The surgeon indicated that the healing abutment may be removed after four weeks. By then, the temporary screw-retained crown had already been fabricated by the technician, who had prepared a CAD/CAM-milled acrylic tooth glued on to a Neoss NeoLink abutment (Figs. 20 & 21). As a result of the decision to place the implant entirely in native bone, the angulation was such as to locate the screw access hole of the provisional on the buccal aspect. This can be easily camouflaged by a simple composite filling after plugging the channel with PTFE tape. The gingival profile copies in this first stage of loading the central incisor anatomy of the Neoss Esthetic Healing Abutment (Fig. 22).

### Tissue conditioning

As evidenced by the aesthetic analysis before treatment, it was clear that symmetry with the contralateral incisor would be impossible. The implant was placed slightly distal because the distal papilla normally has a narrower mesiodistal basis than the central papilla. The tissue volume augmentation helped to obtain the necessary quantity of gingiva to shape nice papillae, leaving a minimal gap. The soft tissue architecture was conditioned by adding composite to the temporary crown and grinding material where necessary until the prosthodontist and the patient felt an optimal result had been achieved.

### Transfer of the profile

Another intra-oral scan of the full arch with the temporary crown in place was performed. The provisional was then removed from the mouth and screwed on to an implant replica fixed to a stable support with wax. A second scan



# Implant dentistry from A to X

Women and men, young and old, that have lost their teeth place their trust in you to restore their happiness and quality of life. It is your detailed planning, careful surgery, unique restorations and individual follow-ups that let them experience the confidence and joy of smiling again.

So whatever challenges you face, you can count on us to develop solutions, concepts and processes that help you in your daily work, help you build your practice and your team so that you can achieve the best results for your patients.

Making the difference together.  
Dentsply Sirona Implants



Comprehensive solutions




Professional and business development



Digital planning



Regenerative



Implant systems



Restorative

Follow Dentsply Sirona Implants



[dentsplysirona.com/implants](https://dentsplysirona.com/implants)








Fig. 26



Fig. 27

revealed in 360° the modified shape of the temporary crown, including the gingival profile (Fig. 23). These files can be easily matched in the CAD software when designing the definitive crown (Figs. 24 & 25).

### Definitive crown

Owing to the angulation of the implant, it was necessary to relocate the screw access hole. In CAD, the design for a cobalt-chromium support that copied the gingival profile of the temporary was prepared, and the screw access was brought to the palatal aspect (Fig. 26). The file was sent to a milling centre in Sweden. High-quality material and CAM production guarantee an excellent outcome in terms of connection and smooth surfaces (Fig. 27). Feldspathic ceramics are layered to obtain the final anatomy and texture. The patient was satisfied with the result. Minor gingival asymmetries are not really disturbing when viewed at social distance if other parameters, like colour, incisal edge, and tooth texture are respected (Fig. 28).



Fig. 28

### Conclusion

Innovative technologies enable very accurate diagnosis and treatment planning. Affordable high-quality CBCT has profoundly changed our profession. In the current case, the detailed X-Mind trium 3D images allowed for planning and performing implant placement in the optimal mesiodistal position. After collecting all of the necessary information and knowing what technology can provide, it is possible that one team will opt for GBR and monolithic crowns, where another might try to minimise the invasiveness of surgery and employ innovative milling strategies. In the actual challenging buccopalatal dimension, the implant was perfectly planned and guided into the centre of the native bone. GBR was limited to the minimum. The fact that bone volume could be matched with the dental preoperative situation and the CAD virtual wax-up made the whole procedure very predictable. Bone volume, bone quality, extent of GBR indicated and the type of prosthodontic solution were all known before starting treatment thanks to the implant planning with the AIS 3D App software.

*Editorial note: This article was originally published by Dental Tribune International in the 1/2019 issue of CAD/CAM—international magazine of digital dentistry. It is reprinted here (as shortened version with editing changes) with permission. Scan the QR code at the top of the article for the full-length version.*

### about the author



**Dr Jakob Zwaan** graduated from Utrecht University in the Netherlands in 1987. He is a member of Digital Dentistry Society and Nederlandse Vereniging voor Parodontologie (Dutch society of periodontics) and President of the 3D-Simplement Association. He has authored numerous scientific publications and spoken widely at international events. He focuses on implant surgery and digital dentistry, and runs a private practice in Calusco d'Adda in Italy.

### contact

**Dr Jakob Zwaan**  
 studio dentistico dr. Jakob Zwaan  
 Via San Rocco 377  
 24033 Calusco d'Adda, Italy  
 Phone: +39 035798811  
 studiozw@yahoo.it





# The unique Tissue Level Implant with Zirconia Collar



Aesthetic gingival area

Zirconia collar

Gingival integration & Osseointegration

Bone area

Pure Titanium body

## Z1<sup>®</sup> Implant Giving you confidence in implantology.

### Proven Technology

- ✓ 98.6%\* success rate
- ✓ Pure Titanium and Y-TZP Zirconia
- ✓ Suitable for all prosthetic solutions

### Proven clinical outcomes for patient safety

- ✓ Anti-bacterial shield
- ✓ Ideal in fresh extraction sockets
- ✓ Immediate aesthetic result

### Proven economics for a profitable implant practice

- ✓ Reduced chairtime
- ✓ Practice development
- ✓ Patient satisfaction

### Proven surgical protocols for a simplified workflow

- ✓ Only 1 surgery
- ✓ No healing abutment
- ✓ Visibility of the connection



Find us online  
[www.tbr.dental](http://www.tbr.dental)



Unique, like your smile

\*Z1 implants are medical devices of class IIb registered by SUDIMPLANT SAS. Information taken from the data of the Sudeinquiry@ program based on 15,534 patients with Z1 implants from 01/2011 to 01/2016.