

Laser-assisted prophylaxis around zirconia implants

France-based dentist Dr Fabrice Baudot has specialised in periodontics, laser-assisted surgery and implant dentistry and has a particular interest in zirconia implants. In this interview with *laser*, Dr Fabrice Baudot, founding member and scientific leader of the European Academy of Ceramic Implantology (EACim), explains why dental lasers have become an essential part of his daily practice and sheds light on the clinical benefits of combining laser-assisted surgery and ceramic implantology.



Peri-implantitis is an increasing problem in today's dental implantology. It is especially associated with titanium implants. What treatment benefits does the laser offer in this context?

Indeed, peri-implantitis is a real scourge for which we do not yet have good therapeutic solutions. Dr Stefan Renvert, in his 2012 book titled *Peri-implantitis*, said, "As in all pathologies, the best form of treatment is very

often prevention and peri-implantitis is no exception." From this point of view, the Er:YAG laser can provide us with certain solutions for the daily practice. The cause of peri-implantitis might be multifactorial, but the microbiological aspect and the contamination of implant surfaces by biofilms seems to be a key point. With its well-demonstrated bactericidal properties, the Er:YAG laser is an excellent complementary instrument to conventional instrumentation for regulating the development of biofilm around implants and cleaning contaminated implant surfaces. We capitalise on two major effects of the Er:YAG laser in the management of peri-implantitis: firstly, there are photo-ablative effects of the laser, which are used for the micro-ablative cleaning of implant surfaces and the removal of inflammatory tissue. These ablative effects allow us to clean so-called peri-implant wounds; secondly, there are photoacoustic effects, which are based on shock waves generated by the Er:YAG laser emission that agitate the water-based antiseptic irrigation solutions that can be used to clean the peri-implant space.

"Indeed, peri-implantitis is a real scourge for which we do not yet have good therapeutic solutions."

When it comes to oral surgeries, we usually exploit the photo-ablative effects of the Er:YAG laser. If we compare the Er:YAG laser to air polishers, which appear to be the reference for cleaning implant surfaces, the Er:YAG laser shows similar results on rough titanium surfaces, but without leaving powder debris on the surgical site. It can be argued that it is as effective and at the same time cleaner than conventional instruments. As opposed to other lasers, the Er:YAG laser is the only laser that can safely treat hard and soft tissue simultaneously without generating uncontrolled thermal effects. It can particularly be used on bone without the risk of thermal damage and even in confined spaces, as is often the case in dental implantology. In peri-implant maintenance, we exploit the photoacoustic effects, leading to an emulsifying of biofilm without altering the delicate tissue-implant interface. The shock waves generated by Er:YAG radia-

tion with a wavelength of 2,940nm cause a 3D expansion of the irrigation solution, and the agitation effect of the solution destabilises the biofilm in a way well beyond what conventional instrumentation is able to achieve. These repeated preventive measures which are tailored to the patient's individual physiology make it possible to maintain peri-implant homoeostasis without altering the surrounding tissue environment. The treatment protocols are very simple to follow and clinical application only takes a few minutes.

"I use the Er:YAG laser for peri-implant maintenance on a daily basis."

Ceramic implants are not particularly associated with peri-implant inflammation. Are there useful laser applications designed for ceramic implants too? What is possible and what isn't?

I use the Er:YAG laser for peri-implant maintenance on a daily basis. This concept of clinical application is based on the emulsion of biofilm that develops around the trans-

gingival part of implant restorations and even beyond, in cases of peri-implantitis. Dr Kenneth S. Kornman's article "The host response to the microbial challenge in periodontitis: Assembling the players", published in 1997 in *Periodontology 2000*, made me realise the importance of exposing pathogens to defence systems to preserve periodontal and peri-implant homoeostasis. The Er:YAG laser and its photoacoustic effects enable the meeting between the pathogenic agents and the immune defence system, as explained by Kornman in his article. In an excellent journal article ("The effect of material characteristics, of surface topography and of implant components and connections on soft tissue integration: A literature review") published on 1 September 2006 in *Clinical Oral Implants Research*, Belgium-based Prof. Eric Rompen elaborated on the importance of implant materials regarding the nature of tissue attachment. Only titanium and zirconia allow hemidesmosome attachment, creating a kind of barrier against biofilm. Other materials used for transgingival connections, such as gold or fired ceramics, do not favour attachment. A real pocket is formed around this type of connection, which can pose a risk factor for the development of peri-implantitis. Deep laser-assisted peri-implant maintenance is particularly indicated around gold or UCLA-type fired ceramic connections.

AD

CALL FOR PAPERS

on laser-assisted ceramic implantology | laser-assisted peri-implantitis therapy

Join our professional media platform now!



www.oemus.com



Hand in your manuscripts and become part of an internationally networked community!

Your benefits as an author:

- Wide international reach
- Author profile on ZWP online
- Free sample copies
- Offprints on demand

Would you like to publish an article on laser-assisted ceramic implantology or laser-assisted peri-implantitis therapy in one of our international publications?

Contact Mr Georg Isbaner at:

g.isbaner@oemus-media.de

or by phone under: +49 341 48474-123.

We look forward to receiving your manuscripts!

With zirconia and titanium connections, the attachment is of better quality and the use of the Er:YAG laser is less justified when it comes to maintenance. When treating established peri-implantitis, the Er:YAG laser with its micro-ablative properties makes, in my opinion, a real difference compared with conventional instrumentation. Again, it is a very clean and efficient tool. It disinfects the treated sites without altering the adjacent surrounding tissue. Numerous studies have shown its safety on titanium, but there are, as far as I know, very few studies showing the effects of the Er:YAG laser on zirconia im-

“In my opinion, the only laser that is truly effective and safe in peri-implant spaces is the Er:YAG laser.”

plants. I have only *in vitro* experience under an operating microscope of the use of the Er:YAG laser on zirconia implant surfaces (Z-Systems and CERALOG) because I have not had to deal with any cases of peri-implantitis around these implants. According to my observations, the implant surface of zirconia implants appears to be completely inert to Er:YAG radiation and reacts much less than titanium. The organic materials which are present on a ceramic implant surface are literally vaporised owing to the micro-ablative effect of the Er:YAG laser, leaving a perfectly clean and visibly unaltered surface as a result. It therefore appears that the Er:YAG laser is very effective in cleaning ceramic implants.

Are there any material-specific factors that need to be particularly considered when treating ceramics with lasers?

In my opinion, the only laser that is truly effective and safe in peri-implant spaces is the Er:YAG laser. As has been done for titanium implants, the effects of laser radiation should be tested at different power levels, under different circumstances and for different exposure times. Micro-ablative effects on zirconia should be observed by means of a scanning electron microscope (SEM) and possible surface alterations, especially in the microtexture, should be evaluated. One thing that seems fundamental to me is the aspect of the thermal rise of zirconia under the effect of Er:YAG radiation. There is only one *in vitro* SEM study, conducted by Dr Stefan Stübinger in 2008 (“Effect of Er:YAG, CO₂ and diode laser irradiation on surface properties of zirconia endosseous dental implants”; *Lasers in Surgery and Medicine*), which shows and evaluates the effect of radiation from three different lasers on zirconia. The carbon dioxide laser seems to al-

ter the surface of zirconia, and the Er:YAG laser penetrates the surface of zirconia. The diode laser appears to be more suitable, but this study has not been confirmed yet by other studies. The clinical application of the diode laser seems to me, when looking at it *in vitro*, much more problematic than the Er:YAG laser, especially in areas close to the bone as is systematically the case in implantology. The risk of collateral thermal effects especially on the bone is of major concern with diode laser radiation.

Alterations of the implant surface under the effect of laser radiation appear to occur only at powers well beyond those effective in destroying biofilm and organic material. Owing to its peak water absorption, the wavelength at 2,940 nm of the Er:YAG laser is effective at the lowest energy levels and therefore cannot, in clinical use, significantly alter the implant surface. The important thing is to be able to manipulate Er:YAG radiation under visual control at high magnification to optimise power settings at the lowest effective energy levels and at tangential incidence to the surface to limit the concentration of energy transmitted to the implant support and thus limit the potential effects of the radiation on the implant surface to be treated. The clinical experience and mode of use of the laser is fundamental and deserves to be widely studied to confirm the safe clinical efficacy of lasers on zirconia implant surfaces.

about the interviewee


Dr Fabrice Baudot is a French dentist specialised in periodontics and implantology. He currently leads a practice that is specialised in laser-assisted microsurgery. His therapeutic approach is always based on minimally invasive surgery. Dr Baudot is frequently invited to speak at international dental conferences, and he is the author of numerous scientific publications.

contact

Dr Fabrice Baudot

Impasse des Trois Pointes 65
34980 Saint-Gély-du-Fesc, France
Phone: +33 4 99060060
dr.baudot34@orange.fr

Interviewee details



BOOK
YOUR
TICKETS
TODAY

LONDON Dentistry SHOW

COMPLIMENTARY
ENTRANCE TO FMC
SUBSCRIBERS

DATE 15 - 16 JANUARY 2021 | VENUE OLYMPIA CENTRAL LONDON UK

LONDON'S INTERNATIONAL DENTAL SHOW

EXPERIENCE

- 70+ ENGAGING LECTURES OVER TWO DAYS
 - WORLD-CLASS SPEAKERS
- A COMPREHENSIVE RANGE OF CLINICAL AND BUSINESS TOPICS
 - 350+ PREMIER BRANDS EXHIBITING
 - SAVE THOUSANDS ON YOUR PURCHASES
- EXCLUSIVE SHOW OFFERS FROM HUNDREDS OF EXHIBITORS
 - EIGHT LECTURE THEATRES
 - COMPLIMENTARY BEER & BUBBLES SPONSORED BY DÜRR DENTAL
 - UNLIMITED TEA & COFFEE ON BOTH DAYS
 - LONDON'S BIGGEST DENTAL SHOW OF 2020

LECTURE ZONES

AESTHETIC
THEATRE

IDDA DIGITAL
DENTISTRY
THEATRE

IN PARTNERSHIP WITH
IDDA

CLINICAL
EXCELLENCE
THEATRE

RISK
MANAGEMENT
HUB

SPONSORED BY
DDU

COMPLIANCE
HUB

BUSINESS
THEATRE

ENHANCED
CPD
THEATRE

BUSINESS
UPDATE
THEATRE

WITH THANKS TO OUR PARTNERS

DIAMOND SPONSOR

Belmont

CORPORATE PARTNER



straumann group

VOCO
THE DENTALISTS

GOLD SPONSORS

align
invisible | zero

Dentsply
Sirona

DÜRR
DENTAL

HENRY SCHEIN®
DENTAL

ivoclar
vivadent
passion vision innovation

OVER 350
PREMIER
BRANDS
EXHIBITING

TWO
DAY PASS
\$199 USD

ORGANISED BY

FMC

EXHIBITION

DIGITAL PARTNERS

DENTISTRY.co.uk

Dentistry
app

MEDIA PARTNER

Dentistry

EDUCATION PARTNER

Dentistry
ON DEMAND

Book your place at www.londondentistryshow.co.uk today

T +44 (0) 1923 851 777 E info@fmc.co.uk W www.fmc.co.uk

