


Single-tooth restoration in the anterior maxilla


Dr João Pedro Almeida & António Korrodi Ritto, Portugal




01

01
Clinical situation
at the start of
orthodontic
treatment in 2011.


02a–d
CBCT images
showing marginal
bone loss around
tooth #21.



02a



02b



02c



02d

Introduction

Replacing an unsalvageable tooth in the anterior maxilla with a dental implant poses certain challenges clinicians must navigate. Due to the high visibility of this area, using an implant system that can facilitate optimum aesthetics by predictably preserving the hard and soft tissue is crucial. In the clinical case described below, an implant system was used to restore a central incisor that has been proven in independent long-term studies to precisely achieve that.^{1,2}

This two-piece tissue-level implant has a parallel-walled design with a shallow thread and is inserted using moderate torques not exceeding 30 Ncm to avoid compression of the surrounding tissue and thus to allow fast and predictable healing, which is a prerequisite for long-term stability and vitality of bone levels and overall tissue health.


The implant treatment described in this case required a multidisciplinary approach and coordination with the orthodontist, as orthodontic treatment had not been completed at the time of implant placement, as the patient was still wearing retainers.

Initial situation


The female patient, aged 30, was referred by her orthodontist owing to the fractured root of tooth #21. She had started orthodontic treatment in 2011 (Fig. 1). After discoloration of the tooth developed, a CBCT scan was taken (in 2021), and it showed marginal bone resorption and bone loss around the root of the endodontically treated tooth #21 (Figs. 2a–d).




03
Removal of the crown and failing post-and-core restoration.




04
Extraction of the remaining root.



05
Extraction socket filled with particulate xenograft material.



06
Grafted site covered with the collagen membrane.







07
Sutured site.

08
Clinical situation after placement of the provisional restoration.



"A minimally invasive implant placement with a moderate insertion torque of a maximum of 30 Ncm is imperative to ensure minimal bone compression to retain the vitality of the bone after implant insertion, allowing healing to progress undisturbed."



Treatment planning

The treatment would involve extraction of the failing tooth and augmentation of the surrounding hard and soft tissue. This would be followed by placement of a two-piece dental implant in this position after healing.

Surgical procedure

The crown and failing post-and-core restoration were removed (Fig. 3), and the remaining root was extracted (Fig. 4). The extraction socket was then carefully curetted to remove any fibrous tissue and filled with particulate xeno-graft material (Bio-Oss, Geistlich; Fig. 5). The site was covered with a collagen membrane (Bio-Gide, Geistlich; Fig. 6), and a full-thickness palatal pedicle graft was rotated and positioned to assure closure and augmented soft-tissue volume. The site was then sutured (Fig. 7), and a provisional restoration (Maryland bridge) was bonded to the adjacent teeth (Fig. 8).

After a healing period of 6.5 months (Fig. 9), the provisional restoration was removed (Fig. 10). The osteotomy was prepared according to the surgical protocol of the implant manufacturer and a two-piece dental implant (Patent™ Dental Implant System, Zircon Medical Management; 4.5 mm in diameter and 11.0 mm in length) was placed equigingival to an insertion torque of 30 Ncm. The soft tissue was adapted around the implant on the labial aspect as part of a minimally invasive grafting procedure to increase the gingival volume (Figs. 11a & b). The connection of the implant was sealed with a PTFE strip and flowable composite, and a new provisional restoration (Maryland bridge) was bonded to the adjacent teeth, covering but not touching the implant (Fig. 12).

09a-d
CBCT images
after 6.5 months
of healing.

11a + b
Clinical situation
after implant
placement and
minimally invasive
grafting to
increase the
gingival volume.

10
Clinical situation
after removal of
the provisional
restoration.

ISMI

INT. SOCIETY OF METAL FREE IMPLANTOLOGY

The innovative specialist society for modern metal-free implantology

Become a member now!

Benefits



Effective public relations

Benefit from a strong community that elevates the marketing of your practice through effective public relations strategies.

Personal online member profile

ISMI provides a personal profile of all active members on their website—free of charge. In addition, the ISMI patient platform provides important information for patients and features a search tool with which patients can find their perfect dentist.



Discount on congress fees

Come and join us! Attend the Annual Meeting of ISMI in Berlin on 9 and 10 May 2025 and benefit from first-rate continuing education. ISMI members receive a special discount on the participation fee.



Online archive for specialists

Get exclusive access to ISMI's extensive online archive. Discuss all relevant questions regarding metal-free implantology with experts and colleagues from around the world and enjoy free access to the online archive where you will find informative training videos and clinical case reports.




Newsletter

The ISMI newsletter keeps you up to date with the latest scientific trends, products, and events on a regular basis. It also features user reports as well as a wide range of information and tips on the subject of metal-free implantology.



Specialist magazine

As a member of ISMI, your membership fee includes a subscription of the independently published English language magazine *ceramic implants—international magazine of ceramic implant technology*. Published three times per year, the magazine offers specialist articles and event reports as well as industry- and science-related news from the international world of metal-free implantology. In addition, *ceramic implants* provides information about manufacturers and their latest products.



ISMI e.V. Office Leipzig
Holbeinstraße 29 | 04229 Leipzig | Germany
Phone: +49 800 4764-000 | Fax: +49 800 4764-100
office@ismi.me | www.ismi.me


Become a
member of ISMI now!





12
Clinical situation
after placement
of the new
provisional
restoration.

13a-c
Clinical situation
and CBCT images
after a further
healing period of
four months.



14
Removal of
excess gingiva.


15
Cementation
of the glass fiber
post.

16
Clinical situation
after intra-oral
preparation of
the glass fiber
post.

17
Clinical situation
after placement
of another
provisional
restoration.



18a + b
Clinical situation
one month later,
after orthodontic
space closure.



19
Fabrication of the
final crown.



Prosthetic restoration

After a further healing period of four months, the patient returned for prosthetic restoration, showing a successfully osseointegrated implant with healthy soft tissue and a stable marginal bone level (Figs. 13a–c). Excess gingiva was removed with an electrosurgical device to expose the margin of the implant (Fig. 14). The glass fibre post of the two-piece implant system used was then cemented into the implant's prosthetic connection using a dual-polymerising dental cement (ACTIVA BioACTIVE-CEMENT, Pulpdent; Fig. 15) and prepared using a diamond bur at high speed under water irrigation (Fig. 16). The prepared post received a provisional crown (Fig. 17), and the patient was sent to the referring orthodontist for space closure before the placement of the final crown.

One month later, the spaces on either side of tooth #21 had been closed (Figs. 18a & b), and a final digital impression was taken. The Matisse-protocol was used to match the shade and achieve the proper colour. The scan files were transferred to the dental laboratory (Dentalook), where the final crown was fabricated (Fig. 19). The patient then received a second provisional crown (Fig. 20). Three months later, the patient received the final crown. The result, two weeks after placement of the final restoration, was deemed highly satisfactory (Fig. 21). At a follow-up eight months later, the soft tissue had matured and was deemed healthy and stable (Figs. 22a & b).

Discussion

The challenges of this clinical case included the lack of marginal bone crest because of external root resorption and the absence of an alveolar buccal wall, necessitating augmentation. Also, since the procedure involved restoring the aesthetic zone, a highly visible area, a dental implant was needed capable of maintaining vital and stable hard and soft tissue. The Patent™ system has been demonstrated to have this capability in long-term studies.^{1,2} These studies have reported healthy soft tissue, an aesthetic increase in keratinised gingiva as well as minimal marginal bone losses after nine years of function, and no peri-implantitis even up to 12 years of function. These results reflect the treatment outcome of the case discussed here, where the



20
Clinical situation
after placement
of another
provisional
restoration.

21
Satisfactory result
two weeks after
placement of the
final crown.

22a + b
Healthy and
stable soft tissue
eight months
later.



patient presented with stable bone levels around her new implant at the follow-up over 17 months after implant placement, as well as healthy soft tissue (Figs. 22a & b).


A minimally invasive implant placement with a moderate insertion torque of a maximum of 30 Ncm is imperative to ensure minimal bone compression to retain the vitality of the bone after implant insertion, allowing healing to progress undisturbed. This is crucial to maintaining long-term stability of the surrounding tissue. We have found adopting such a low torque strategy together with an atraumatic insertion protocol with this implant system to minimise marginal bone loss and to maintain overall tissue stability in daily practice.

Conclusion

The implant system used in this clinical case represents a viable option for replacing teeth in the aesthetic zone. Owing to the implant's ability to maintain vital and stable hard and soft tissue and to foster strong soft-tissue adhesion, highly satisfactory aesthetic results are to be expected over the long term, and the risk of bacteria-induced chronic tissue inflammation like peri-implantitis is minimised.^{1,2} The healing of the soft tissue in this case progressed rapidly with almost no inflammation.

Literature:

- ¹ Brunello G, Rauch N, Becker K, Hakimi AR, Schwarz F, Becker J. Two-piece zirconia implants in the posterior mandible and maxilla: a cohort study with a follow-up period of 9 years. Clin Oral Implants Res. 2022. december;33(12):1233–44. doi: 10.1111/clr.14005. PMID: 36184914.
- ² Karapataki S, Vegh D, Payer M, Fahrenholz H, Antonoglou GN. Clinical performance of two-piece zirconia dental implants after 5 and up to 12 years. Int J Oral Maxillofac Implants. 2023. december 12.;38(6):1105–14. doi: 10.11607/jomi.10284. PMID: 38085741.



Dr João Pedro Almeida

Clínica Médica de Implantologia

R. Comissão de Iniciativa

Nº2 A, Piso 6

2410-098 Leiria

Portugal


+34 244 824436

www.clinicaimplantologia.pt

Dr João Pedro
Almeida



Literature



ZIRKONUS

Implantatsysteme



Featuring a patented ceramic-compatible coupling mechanism for seamless integration between implant, customised abutment, and crown. Available in single-piece and multi-piece options, with diameters from 3.5 to 11.5 mm and lengths from 6 to 14 mm.

ZIRKONUS Implantatsysteme GmbH & Co. KG

Bahnhofstraße 18, 71034 Böblingen, Germany
☎ +49 711 305329-14

info@zirkonus.de
www.zirkonus.de

CE 0483

